arXiv:1301.3404v1 [math.PR] 15 Jan 2013

Pólya urns via the contraction method

Margarete Knape and Ralph Neininger
Institute for Mathematics
J.W. Goethe University
60054 Frankfurt a.M.
Germany

Email: {knape,neiningr}@math.uni-frankfurt.de

January 15, 2013

Abstract

We propose an approach to analyze the asymptotic behavior of Pólya urns based on the contraction method. For this a combinatorial discrete time embedding of the evolution of the composition of the urn into random rooted trees is used. A decomposition of the trees leads to a system of recursive distributional equations which capture the distributions of the numbers of balls of each color. Ideas from the contraction method are used to study such systems of recursive distributional equations asymptotically. We apply our approach to a couple of concrete Pólya urns that lead to limit laws with normal limit distributions, with non-normal limit distributions and with asymptotic periodic distributional behavior.

MSC2010: 60C05, 60F05, 60J05, 68Q25.
Keywords: Pólya urn, Pólya-Eggenberger urn, contraction method, weak convergence, limit law, probability metric, recursive distributional equation.

1 Introduction

In this paper we develop an approach to prove limit laws for Pólya urn models by the contraction method. We consider an urn with balls of a finite number $m \geq 2$ of different colors, numbered by $1, \ldots, m$. The evolution of a Pólya urn is determined by an $m \times m$ replacement matrix $R = (a_{ij})_{1 \leq i,j \leq m}$ which is given in advance together with an initial (time 0) composition of the urn with at least one ball. Time evolves in discrete steps. In each step one ball is drawn uniformly at random from the urn. If it has color i it is placed back to the urn together with a_{ij} balls of color j for all $j = 1, \ldots, m$. The steps are iterated independently. A classical problem is to identify the asymptotic behavior of the numbers of balls of each color as the number n of steps tends to infinity. The literature on this problem, in particular on limit laws for the normalized numbers of balls of each color, is vast. We refer to the monographs of Johnson and Kotz [14] and Mahmoud [18] and the references and comments on the literature in the papers of Janson [12], Flajolet et al. [10] and Pouyanne [23].

A couple of approaches have been used to analyze the asymptotic behavior of Pólya urn models, most notably the method of moments, discrete time martingale methods, embeddings into continuous time multitype branching processes, and methods from analytic combinatorics based on generating functions. All these methods use the “forward” dynamic of the urn process by exploiting that the distribution of the composition at time n given time $n-1$ is explicitly accessible.

In the present paper we propose an approach based on a “backward” decomposition of the urn process. We construct an embedding of the evolution of the urn into an associated combinatorial random tree structure growing in discrete time. Our associated tree can be decomposed at its root (time 0) such that the growth dynamics of the subtrees of the root resemble the whole tree in distribution. More precisely we have different types of distributions for the associated tree, one type for each possible color of its root. The decomposition of the associated tree into subtrees gives rise to a system of distributional recurrences for the numbers of balls of each color. To extract the asymptotic behavior from such systems we develop an approach in the context of the contraction method.

The contraction method is well known in the probabilistic analysis of algorithms. It was introduced by Rösler [25] and first developed systematically in Rachev and Rüschendorf [24]. A rather general framework with numerous applications to the analysis of recursive algorithms and random trees was given by Neininger and Rüschendorf [20]. The contraction method has been used for sequences of distributions of random variables (or random vectors or stochastic processes) that satisfy an appropriate recurrence relation. To the best of our knowledge it has not yet been used for systems of such recurrence relations as they arise in the present paper, the only exception being Leckey et al. [17] where tries are analyzed under a Markov source model.

The aim of this paper is not to compete with other techniques with respect to generality under which urn models can be analyzed. Instead we discuss our approach at a couple of examples illustrating the contraction framework in three frequently occurring asymptotic regimes: normal limit laws, non-normal limit laws and regimes with oscillating distributional behavior. We also discuss the case of random entries in the replacement matrix. Our proofs are generic and can easily be transfered to other urn models or be developed into more general theorems when asymptotic expansions of means (respectively means and variances in the normal limit case) are available, cf. the types of expansions of the means in section 3.

A general assumption in the present paper is that the replacement matrix is balanced, i.e., that we have $\sum_{j=1}^m a_{ij} =: K-1$ for all $i = 1, \ldots, m$, where $K \geq 2$ is a fixed integer. (The notation K is unfortunate since this integer is not random and mainly chosen to have similarity in notation with earlier work on the contraction method.) An implication of the balance condition is that the growths of the subtrees of the associated tree processes can asymptotically jointly be captured by Dirichlet distributions. This leads to characterizations of the limit distributions in all cases (normal, non-normal and oscillatory limits) by systems, cf. (6)–(8) below, of distributional fixed-point equations where all coefficients are powers of components of a Dirichlet distributed vector, see also the discussion in section 3. It may be an interesting aspect of the present approach that all three regimes are governed by these quite similar types of systems of distributional fixed-point equations.

The paper is organized as follows: in section 2 we introduce the associated trees into which the urn models are embedded and derive the systems of distributional recurrences for the numbers of balls of a certain color from the associated trees. In section 3 we outline the types of systems of fixed-point equations that emerge from the distributional recurrences after a proper normalization. To make these recurrences and fixed-point equations accessible to the contraction method, in section 4 we first introduce spaces of distributions and appropriate cartesian product spaces together with metrics on these product spaces. The metrics in use are product versions of the minimal L_p metrics and product versions of the Zolotarev metrics. In section 5 we use these spaces and metrics to show that our systems of distributional fixed-point equations uniquely characterize vectors of probability distributions via a contraction property. These cover the types of distributional fixed-point equations that appear in the final section 6 where we discuss examples of limit laws for Pólya urn schemes within our approach. In section 6 also our convergence proofs are worked out, again based on the product versions of the minimal L_p and Zolotarev metrics.

Notation. By $\xrightarrow{d}$ convergence in distribution is denoted. We denote the normal distribution on $\mathbb{R}$ with mean $\mu \in \mathbb{R}$ and variance $\sigma^2 \geq 0$ by $\mathcal{N}(\mu, \sigma^2)$. In the case $\sigma^2 = 0$ this degenerates to the Dirac measure in 0. Throughout the Bachmann-Landau symbols are used in asymptotic statements. By $\log(x)$ for $x > 0$ the natural logarithm of x is denoted.

2 A recursive description of Pólya urns

In this section we explain the embedding of urn processes into associated combinatorial random tree structures growing in discrete time. The distributional self-similarity within the subtrees of the roots of these associated trees leads to systems of distributional recurrences which are the basis of our approach.

The Pólya urn. To develop our approach we first consider an urn model with two colors, black and white, and a deterministic replacement matrix R, where the sums of the entries in each row is constant. Below, an extension of this approach to urns with more than two colors and replacement matrices with random entries is discussed as well. However, the assumption that the sums of the entries in each row is the same is crucial for our method, the reason also being explained below. To be definite we use the replacement matrix

$$R = \begin{bmatrix} a & b \\ c & d \end{bmatrix} \qquad \text{with } a, d \in \mathbb{N}_0 \cup \{-1\} \text{ and } b, c \in \mathbb{N}_0 \tag{1}$$

with

$$a + b = c + d =: K - 1 \geq 1.$$

Hence after drawing a black ball this ball is placed back to the urn together with a new black balls and b new white balls. If drawing a white ball it is placed back to the urn together with c black balls and d white balls. A diagonal entry $a = -1$ (or $d = -1$) implies that a drawn black (or white) ball is not replaced to the urn while balls of the other color are still added to the urn. As initial configuration we consider both, one black ball or one white ball. Other initial configurations can be dealt with as well, also discussed below. We denote by B_n^{b} the number of black balls after n steps when initially starting with one black ball, by B_n^{w} the number of black balls after n steps when initially starting with one white ball. Hence we have $B_0^{\mathrm{b}} = 1$ and $B_0^{\mathrm{w}} = 0$.

The associated tree. We encode the urn process as follows by a discrete time evolution of a random tree with nodes colored black or white. This tree is called *associated tree*. The initial urn with one ball, say a black one, is associated with a tree with one root node of the same (black) color. The ball in the urn is represented by this root node. Now drawing the ball and placing it back to the urn together with a new black balls and b new white balls is encoded in the associated tree by adding $a+b+1=K$ children to the root node, $a+1$ of them being black and b being white. The root node then no longer represents a ball in the tree, whereas the K new leaves of the tree now represent the K balls in the urn. Now, we iterate this procedure: At any step, a ball is drawn from the urn. It is represented by one of the leaves, say node v in the tree. The urn follows its dynamic. If the ball drawn is black, the (black) leaf v gets K children, $a+1$ black ones and b white ones. Similarly, if the ball drawn is white, the (white) leaf v gets c black children and $d+1$ white children. In both cases v no longer represents a ball in the urn. The ball drawn and the new balls are represented by the children of v. The correspondence between all other leaves of the tree and the other balls in the urn remains unchanged. For an example of an evolution of an urn and its associated tree see Figure 1. Hence, at any time the balls in the urn are represented by the leaves of the associate tree, where the colors of balls and representing leaves match. Each node of the tree is either a leaf or has K children. We could as well simulate the urn process by only running the evolution of the associated tree as follows: Start with one root node of the color of the initial ball of the urn. At any step choose one of the leaves of the tree uniformly at random, inspect its color, add K children to the chosen leaf and color these children as defined above. Then after n steps the tree has $n(K-1)+1$ leaves, the number of black leaves is distributed as B_n^{b} if the root node was black and distributed as B_n^{w}, if the root node was white.



Figure 1: A realization of the evolution of the Pólya urn with replacement matrix $\left[\begin{smallmatrix}1 & 2\\ 2 & 1\end{smallmatrix}\right]$ and initially one white ball. The arrows indicate which ball is drawn (resp. node is replaced) in each step. Below each urn its associated tree is shown. Leaf nodes correspond to the balls in the urn, non-leaf nodes (crossed out) do no longer correspond to balls in the urn. However, their color still matters for the recursive decomposition of the associated tree.

Subsequently it is important to note the following recursive structure of the associated tree: For a fixed replacement matrix of the Pólya urn we consider the two initial compositions of one black respectively one white ball and their two associated trees. We call these the b-associated respectively w-associated tree. Consider one of these associated trees after n steps. It has $n(K-1)+1$ leaves and each subtree rooted at a child of the associated tree's root (we call them shortly only subtrees) has a random number of leaves according to how often a leaf node has been chosen for replacement in the subtree. We condition on the numbers of leaves of the subtrees to be $i_r(K-1)+1$ with $i_r \in \mathbb{N}_0$ for $r = 1, \dots, K$. Note that we have $\sum_{r=1}^{K} i_r = n-1$, the -1 resulting from the fact that in the first step of the evolution of the associated tree the subtrees are being generated, only afterwards they start growing. From the evolution of the b-associated tree it is clear that conditionally on the subtrees' numbers of leaves being $i_r(K-1)+1$ the subtrees are stochastically independent and the r-th subtree is distributed as an associated tree after i_r steps. Whether it has the distribution of the b- or the w-associated tree depends on the color of the subtree's root node.

To summarize we have that conditionally on their numbers of leaves the subtrees of associated trees are independent and distributed as associated trees of corresponding size and type inherited from the color of their root node.

Growth of subtrees. In our analysis the asymptotic growth of the K subtrees of the associated tree is used. We denote by $I^{(n)} = (I_1^{(n)}, \dots, I_K^{(n)})$ the vector of the numbers of draws of leaves from each subtree after n draws in the full associated tree. In other words, $I_r^{(n)}(K-1)+1$ is the number of leaves of the r-th subtree after $n \in \mathbb{N}_0$ steps. We have $I^{(1)} = (0, \dots, 0)$, and $I^{(2)}$ is a vector with all entries being 0 except for one coordinate which is 1. To describe the asymptotic growth of $I^{(n)}$ we need the Dirichlet distribution $\mathrm{Dirichlet}\big((K-1)^{-1}, \dots, (K-1)^{-1}\big)$: It is the distribution of a random vector $(D_1, \dots, D_K)$ with $\sum_{r=1}^{K} D_j = 1$ and such that $(D_1, \dots, D_{K-1})$ has a Lebesgue-density supported by the simplex $\mathcal{S}_K := \{(x_1, \dots, x_{K-1}) \in [0,1]^{K-1} \mid \sum_{r=1}^{K-1} x_r \le 1\}$ given by

$$x = (x_1, \dots, x_{K-1}) \mapsto c_K \prod_{r=1}^{K-1} x_r^{\frac{2-K}{K-1}}, \quad x \in \mathcal{S}_K, \qquad c_K = \frac{\Gamma((K-1)^{-1})^{1-K}}{K-1},$$

where Γ denotes the gamma function. In particular, $D_1, \dots, D_K$ are identically distributed with the $\mathrm{beta}((K-1)^{-1}, 1)$ distribution, i.e., with Lebesgue-density

$$x \mapsto (K-1)x^{\frac{2-K}{K-1}}, \quad x \in [0,1].$$

We have the following asymptotic behavior of $I^{(n)}$:

Lemma 2.1. *Consider a Pólya urn with constant row sum $K-1 \ge 1$ and its associated tree. For the numbers of balls $I^{(n)} = (I_1^{(n)}, \dots, I_K^{(n)})$ drawn in each subtree of the associated tree when n balls have been drawn in the whole associated tree we have, as $n \to \infty$,*

$$\left(\frac{I_1^{(n)}}{n}, \dots, \frac{I_K^{(n)}}{n}\right) \longrightarrow (D_1, \dots, D_K)$$

almost surely and in any L_p, where $(D_1, \dots, D_K)$ has the Dirichlet distribution

$$\mathcal{L}(D_1, \dots, D_K) = \mathrm{Dirichlet}\left(\frac{1}{K-1}, \dots, \frac{1}{K-1}\right).$$

Proof. The sequence $(I_1^{(n)}(K-1)+1,\dots,I_K^{(n)}(K-1)+1)_{n\in\mathbb{N}_0}$ has an interpretation by another urn model, which we call the subtree-induced urn: For this we give additional labels to the leaves of the associated tree. The set of possible labels is $\{1,\dots,K\}$ and we label a leaf j if it belongs to the j-th subtree of the root (any ordering of the subtrees of the root is fine). Hence, all leaves of a subtree of the associated tree's root get the same label, leaves of different subtrees get different labels. Now, the subtree-induced urn has balls of colors $1,\dots,K$, at each time the number of balls of each color is identical with the numbers of leaves of the corresponding label. Hence the dynamic of the subtree-induced urn is that of a Pólya urn with initially K balls, one of each color. Whenever a ball is drawn, it is placed back to the urn together with $K-1$ balls of the same color. In other words the replacement matrix for the dynamic of the subtree-induced urn is a $K\times K$ diagonal matrix with all diagonal entries being $K-1$. After n steps, we have $I_r^{(n)}(K-1)+1$ balls of color r. The dynamic of the subtree-induced urn as a multivariate Pólya-Eggenberger urn is well-known, we have

$$\left(\frac{I_1^{(n)}(K-1)+1}{n(K-1)+1},\dots,\frac{I_K^{(n)}(K-1)+1}{n(K-1)+1}\right)\longrightarrow(D_1,\dots,D_K)$$

almost surely as $n\to\infty$ where $(D_1,\dots,D_K)$ has the Dirichlet$((K-1)^{-1},\dots,(K-1)^{-1})$ distribution. This implies the assertion. ☐

Note that a Pólya urn violating the condition of constant row sums in the replacement matrix has an associated tree with an $I^{(n)}$ such that $I^{(n)}/n$ converges to a degenerate limit, a deterministic vector with one 1 and all other components 0. However, a non-degenerate limit for $I^{(n)}/n$ will be crucial below.

System of recursive equations. We set up recursive equations for the distributions of the quantities B_n^{b} and B_n^{w}: For B_n^{b} we start the urn with one black ball and have a b-associated tree with a black root node. Now, B_n^{b} is distributed as the number of black leaves in the associated tree after n steps which we express as the sum of the numbers of black leaves of its subtrees. As discussed above, conditionally on $I^{(n)}=(I_1^{(n)},\dots,I_K^{(n)})$, the vector of the numbers of balls drawn in each subtree, these subtrees are independent and distributed as b-associated trees or w-associated trees of the corresponding size depending on the color of their roots. A b-associated tree has $a+1$ black and $b=K-(a+1)$ white children. Hence we obtain

$$B_n^{\mathrm{b}}\stackrel{d}{=}\sum_{r=1}^{a+1}B_{I_r^{(n)}}^{\mathrm{b},(r)}+\sum_{r=a+2}^{K}B_{I_r^{(n)}}^{\mathrm{w},(r)},\quad n\ge 1,\tag{2}$$

where $\stackrel{d}{=}$ denotes that left and right hand side have identical distribution, we have that $(B_k^{\mathrm{b},(1)})_{0\le k<n},\dots,(B_k^{\mathrm{b},(a+1)})_{0\le k<n},(B_k^{\mathrm{w},(a+2)})_{0\le k<n},\dots,(B_k^{\mathrm{w},(K)})_{0\le k<n},I^{(n)}$ are independent, the $B_k^{\mathrm{b},(r)}$ are distributed as B_k^{b}, the $B_k^{\mathrm{w},(r)}$ are distributed as B_k^{w} for $k=0,\dots,n-1$ and the respective values of r.

Similarly, we obtain a recursive distributional equation for B_n^{w}. We have

$$B_n^{\mathrm{w}}\stackrel{d}{=}\sum_{r=1}^{c}B_{I_r^{(n)}}^{\mathrm{b},(r)}+\sum_{r=c+1}^{K}B_{I_r^{(n)}}^{\mathrm{w},(r)},\quad n\ge 1,\tag{3}$$

with conditions on independence and identical distributions as in (2). Note that with the initial value $(B_0^{\mathrm{b}}, B_0^{\mathrm{w}}) = (1,0)$ the system of equations (2)–(3) defines the sequence of pairs of distributions $(\mathcal{L}(B_n^{\mathrm{b}}), \mathcal{L}(B_n^{\mathrm{w}}))_{n\geq 0}$.

General number of colors. The approach above for urns with two colors extends directly to urns with an arbitrary number $m \geq 2$ of colors. We denote the replacement matrix by $R = (a_{ij})_{1\leq i,j\leq m}$ with

$$a_{ij} \in \begin{cases} \mathbb{N}_0, & \text{for } i \neq j, \\ \mathbb{N}_0 \cup \{-1\}, & \text{for } i = j, \end{cases} \qquad \text{and} \qquad \sum_{j=1}^{m} a_{ij} =: K-1 \geq 1 \text{ for } i = 1, \ldots, m.$$

The colors (subsequently also called types) are now numbered $1, \ldots, m$ and we focus on the number of balls of color 1 after n steps. When starting with one ball of color j we denote by $B_n^{[j]}$ the number of color 1 balls after n steps. To formulate a system of distributional recurrences generalizing (2) and (3) we further denote the intervals of integers

$$J_{ij} := \begin{cases} \left[1 + \sum_{k<i} a_{kj}, \sum_{k\leq i} a_{kj}\right] \cap \mathbb{N}_0, & \text{for } i < j, \\ \left[1 + \sum_{k<i} a_{kj}, 1 + \sum_{k\leq i} a_{kj}\right] \cap \mathbb{N}_0, & \text{for } i = j, \\ \left[2 + \sum_{k<i} a_{kj}, 1 + \sum_{k\leq i} a_{kj}\right] \cap \mathbb{N}_0, & \text{for } i > j, \end{cases} \tag{4}$$

with the convention $[x,y] = \emptyset$ if $x > y$. Then, we have

$$B_n^{[j]} \stackrel{d}{=} \sum_{i=1}^{m} \sum_{r\in J_{ij}} B_{I_r^{(n)}}^{[i],(r)}, \quad n \geq 1, \quad j \in \{1, \ldots, m\}, \tag{5}$$

where, for each $j \in \{1, \ldots, m\}$ we have that the family

$$\left\{ \left(B_k^{[i],(r)}\right)_{0\leq k<n} \,\middle|\, r \in J_{ij}, i \in \{1, \ldots, m\} \right\} \cup \left\{ I^{(n)} \right\}$$

is independent, $B_k^{[i],(r)}$ is distributed as $B_k^{[i]}$ for all $i \in \{1, \ldots, m\}$, $0 \leq k < n$ and $r \in J_{ij}$ and $I^{(n)}$ has the distribution as above in Lemma 2.1.

Random entries in the replacement matrix. The case of a replacement matrix with random entries such that all rows sum up to a deterministic and fixed $K-1 \geq 1$ can be covered by an extension of the system (5). Instead of formulating such an extension explicitly, we discuss an example in section 6.2.

3 Systems of limit equations

We outline how systems of the form (5) are used subsequently. Crucial are the expansions of the means

$$\mu_n^{[j]} := \mathbb{E}\left[B_n^{[j]}\right], \quad j = 1, \ldots, m.$$

We only consider cases where these means grow linearly. Note however, that even balanced urns can have quite different growth orders, an example is the replacement matrix $\left[\begin{smallmatrix} 4 & 0 \\ 3 & 1 \end{smallmatrix}\right]$, see Kotz et al. [16].

Type (a). Assume that we have expansions of the form, as $n \to \infty$,

$$\mu_n^{[j]} = c_\mu n + d_j n^\lambda + o(n^\lambda), \quad j = 1, \ldots, m,$$

with a constant $c_\mu > 0$ independent of j, with constants $d_j \in \mathbb{R}$ and an exponent $1/2 < \lambda < 1$. We call this scenario of **type (a)**. This suggests that the variances are of the order $n^{2\lambda}$ and a proper scaling is

$$X_n^{[j]} := \frac{B_n^{[j]} - \mu_n^{[j]}}{n^\lambda}, \quad n \geq 1, \quad j = 1, \ldots, m.$$

Deriving from (5) a system of recurrences for the $X_n^{[j]}$ and letting formally $n \to \infty$ (this is done explicitly in the examples in section 6) we obtain the system of fixed-point equations

$$X^{[j]} \stackrel{d}{=} \sum_{i=1}^{m} \sum_{r \in J_{ij}} D_r^\lambda X^{[i],(r)} + b^{[j]}, \quad j = 1, \ldots, m, \tag{6}$$

where the $X^{[i],(r)}$ and $(D_1, \ldots, D_K)$ are independent, the $X^{[i],(r)}$ are distributed as $X^{[i]}$, $(D_1, \ldots, D_K)$ is distributed as in Lemma 2.1 and the $b^{[j]}$ are functions of $(D_1, \ldots, D_K)$. It turns out that such a system subject to centered $X^{[j]}$ with finite second moments has a unique solution on the level of distributions (Theorem 5.1). This identifies the weak limits of the $X_n^{[j]}$. Examples are in sections 6.1 and 6.2. One can as well obtain the same system (6) with $b^{[j]} = 0$ for all j by only centering the $B_n^{[j]}$ by $c_\mu n$ instead of the exact mean. Then, system (6) has to be solved subject to finite second moments and appropriate means.

Type (b). Assume that we have expansions of the form, as $n \to \infty$,

$$\mu_n^{[j]} = c_\mu n + o(\sqrt{n}), \quad j = 1, \ldots, m,$$

with a constant $c_\mu > 0$ independent of j. We call this scenario of **type (b)**. This suggests that the variances are of linear order and a proper scaling is

$$X_n^{[j]} := \frac{B_n^{[j]} - \mu_n^{[j]}}{\sqrt{\mathrm{Var}(B_n^{[j]})}}, \quad n \geq 1, \quad j = 1, \ldots, m$$

(or $\sqrt{\mathrm{Var}(B_n^{[j]})}$ replaced by $\sqrt{n}$). The corresponding system of fixed-point equations in the limit is

$$X^{[j]} \stackrel{d}{=} \sum_{i=1}^{m} \sum_{r \in J_{ij}} \sqrt{D_r} X^{[i],(r)}, \quad j = 1, \ldots, m, \tag{7}$$

with conditions as in (6). Under appropriate assumptions on moments we find that the only solution is all $X^{[j]}$ being standard normally distributed (Theorem 5.2). This leads to asymptotic normality of the $X_n^{[j]}$. Examples are given in sections 6.1 and 6.2. The case

$$\mu_n^{[j]} = c_\mu n + \Theta(\sqrt{n}), \quad j = 1, \ldots, m,$$

leads to the same system of fixed-point equations (7). However, here the variances typically are of order $n\log^{\delta}(n)$ with a positive δ.

Type (c). Assume that we have expansions of the form, as $n \to \infty$,

$$\mu_n^{[j]} = c_\mu n + \Re\left(\kappa_j n^{\mathrm{i}\mu}\right) n^{\lambda} + o(n^{\lambda}), \quad j = 1, \ldots, m,$$

with a constant $c_\mu > 0$ independent of j, $1/2 < \lambda < 1$, constants $\kappa_j \in \mathbb{C}$ and $\mu \in \mathbb{R}$. (By i the imaginary unit is denoted.) We call this scenario of **type (c)**. This suggests oscillating variances of the order $n^{2\lambda}$. The oscillatory behavior of mean and variance can typically not be removed by proper scaling to obtain convergence towards a fixed limit distribution. Using the scaling

$$X_n^{[j]} := \frac{B_n^{[j]} - c_\mu n}{n^{\lambda}}, \quad n \geq 1, \quad j = 1, \ldots, m.$$

it turns out that the oscillating behavior of the $X_n^{[j]}$ can be captured by the system of fixed-point equations

$$X^{[j]} \stackrel{d}{=} \sum_{i=1}^{m} \sum_{r \in J_{ij}} D_r^{\omega} X^{[i],(r)}, \quad j = 1, \ldots, m, \tag{8}$$

with conditions as in (6) and $\omega := \lambda + \mathrm{i}\mu$. Under appropriate moment assumptions this has a unique solution within distributions on $\mathbb{C}$ (Theorem 5.3). An example of a corresponding distributional approximation is given in section 6.3.

Note that the approach of embedding urn models into continuous time multitype branching processes, see [1, 12], also leads to characterizations of the non-normal limits as in (6) and (8). However, the form of the fixed-point equations is different, see the system in equation (3.5) in Janson [12]. Properties of such fixed-points have been studied in Chauvin et al. [6, 7].

4 Spaces of distributions and metrics

In this section we define cartesian products of spaces of probability distributions and metrics on these products. These metric spaces will be used below to first characterize limit distributions of urn models (section 5) and then prove convergence in distribution of the scaled numbers of balls of a color (section 6).

Spaces. We denote by $\mathcal{M}^{\mathbb{R}}$ the space of all probability distributions on $\mathbb{R}$ with the Borel σ-field. Moreover, we consider the subspaces

$$\mathcal{M}_s^{\mathbb{R}} := \left\{ \mathcal{L}(X) \in \mathcal{M}^{\mathbb{R}} \,\middle|\, \mathbb{E}[|X|^s] < \infty \right\}, \quad s > 0, \tag{9}$$

$$\mathcal{M}_s^{\mathbb{R}}(\mu) := \left\{ \mathcal{L}(X) \in \mathcal{M}_s^{\mathbb{R}} \,\middle|\, \mathbb{E}[X] = \mu \right\}, \quad s \geq 1, \mu \in \mathbb{R} \tag{10}$$

$$\mathcal{M}_s^{\mathbb{R}}(\mu, \sigma^2) := \left\{ \mathcal{L}(X) \in \mathcal{M}_s^{\mathbb{R}}(\mu) \,\middle|\, \mathrm{Var}(X) = \sigma^2 \right\}, \quad s \geq 2, \mu \in \mathbb{R}, \sigma \geq 0. \tag{11}$$

We need the d-fold cartesian products, $d \in \mathbb{N}$, of these spaces denoted by

$$\left(\mathcal{M}_s^{\mathbb{R}}\right)^{\times d} := \mathcal{M}_s^{\mathbb{R}} \times \cdots \times \mathcal{M}_s^{\mathbb{R}}, \tag{12}$$

and analogously $(\mathcal{M}_s^{\mathbb{R}}(\mu))^{\times d}$ and $(\mathcal{M}_s^{\mathbb{R}}(\mu, \sigma^2))^{\times d}$.

We also need probability distributions on the complexe plane $\mathbb{C}$. By $\mathcal{M}^{\mathbb{C}}$ the space of all probability distributions on $\mathbb{C}$ with the Borel σ-field is denoted. Moreover, for $\gamma \in \mathbb{C}$ we use the subspaces and product space

$$\begin{aligned}\mathcal{M}_s^{\mathbb{C}} &:= \left\{\mathcal{L}(X) \in \mathcal{M}^{\mathbb{C}} \,\middle|\, \mathbb{E}[|X|^s] < \infty\right\}, \quad s > 0,\\ \mathcal{M}_2^{\mathbb{C}}(\gamma) &:= \left\{\mathcal{L}(X) \in \mathcal{M}_2^{\mathbb{C}} \,\middle|\, \mathbb{E}[X] = \gamma\right\},\\ \left(\mathcal{M}_2^{\mathbb{C}}(\gamma)\right)^{\times d} &:= \mathcal{M}_2^{\mathbb{C}}(\gamma) \times \cdots \times \mathcal{M}_2^{\mathbb{C}}(\gamma).\end{aligned}$$

To cover the different behavior of the urns two types of metrics are constructed, extensions of the Zolotarev metrics ζ_s and the minimal L_p-metric ℓ_p to the product spaces defined above.

Zolotarev metric. The Zolotarev metric has been introduced and studied in Zolotarev [30, 31]. The contraction method based on the Zolotarev metric was systematically developed in [20] and, for issues that go beyond what is needed in this paper, in [15] and [21]. We only need the following properties: For distributions $\mathcal{L}(X)$, $\mathcal{L}(Y) \in \mathcal{M}^{\mathbb{R}}$ the Zolotarev distance ζ_s, $s > 0$, is defined by

$$\zeta_s(X, Y) := \zeta_s(\mathcal{L}(X), \mathcal{L}(Y)) := \sup_{f \in \mathcal{F}_s} |\mathbb{E}[f(X) - f(Y)]| \tag{13}$$

where $s = m + \alpha$ with $0 < \alpha \leq 1$, $m \in \mathbb{N}_0$, and

$$\mathcal{F}_s := \{f \in C^m(\mathbb{R}, \mathbb{R}) : |f^{(m)}(x) - f^{(m)}(y)| \leq |x - y|^\alpha\}, \tag{14}$$

the space of m times continuously differentiable functions from $\mathbb{R}$ to $\mathbb{R}$ such that the m-th derivative is Hölder continuous of order α with Hölder-constant 1.

We have that $\zeta_s(X, Y) < \infty$, if all moments of orders $1, \ldots, m$ of X and Y are equal and if the s-th absolute moments of X and Y are finite. Since later on the cases $1 < s \leq 3$ are used, we have two basic cases: First, for $1 < s \leq 2$ we have $\zeta_s(X, Y) < \infty$ for $\mathcal{L}(X)$, $\mathcal{L}(Y) \in \mathcal{M}_s^{\mathbb{R}}(\mu)$ for any $\mu \in \mathbb{R}$. Second, for $2 < s \leq 3$ we have $\zeta_s(X, Y) < \infty$ for $\mathcal{L}(X)$, $\mathcal{L}(Y) \in \mathcal{M}_s^{\mathbb{R}}(\mu, \sigma^2)$ for any $\mu \in \mathbb{R}$ and $\sigma \geq 0$. Moreover, the pairs $(\mathcal{M}_s^{\mathbb{R}}(\mu), \zeta_s)$ for $1 < s \leq 2$ and $(\mathcal{M}_s^{\mathbb{R}}(\mu, \sigma^2), \zeta_s)$ for $2 < s \leq 3$ are complete metric spaces; for the completeness see [8, Theorem 5.1].

Convergence in ζ_s implies weak convergence on $\mathbb{R}$. Furthermore, ζ_s is $(s, +)$ ideal, i.e., we have

$$\zeta_s(X + Z, Y + Z) \leq \zeta_s(X, Y), \quad \zeta_s(cX, cY) = c^s \zeta_s(X, Y) \tag{15}$$

for all Z being independent of (X, Y) and all $c > 0$. Note, that this implies for $X_1, \ldots, X_n$ independent and $Y_1, \ldots, Y_n$ independent such that the respective ζ_s distances are finite that

$$\zeta_s\left(\sum_{i=1}^n X_i, \sum_{i=1}^n Y_i\right) \leq \sum_{i=1}^n \zeta_s(X_i, Y_i). \tag{16}$$

On the product spaces $(\mathcal{M}_s^{\mathbb{R}}(\mu))^{\times d}$ for $1 < s \leq 2$ and $(\mathcal{M}_s^{\mathbb{R}}(\mu, \sigma^2))^{\times d}$ for $2 < s \leq 3$ our first main tool is

$$\zeta_s^{\vee}((\nu_1, \dots, \nu_d), (\mu_1, \dots, \mu_d)) := \max_{1 \leq j \leq d} \zeta_s(\nu_j, \mu_j), \tag{17}$$

where $(\nu_1, \dots, \nu_d), (\mu_1, \dots, \mu_d) \in \mathcal{M}_s^{\mathbb{R}}(\mu))^{\times d}$ and $\in (\mathcal{M}_s^{\mathbb{R}}(\mu, \sigma^2))^{\times d}$ respectively. Note that $\zeta_s^{\vee}$ is a complete metric on the respective product spaces and induces the product topology.

Minimal $\mathbf{L_p}$-metric $\boldsymbol{\ell_p}$. First for probability metrics on the real line the minimal L_p-metric ℓ_p, $1 \leq p < \infty$ is defined by

$$\ell_p(\nu, \varrho) := \inf\{\|V - W\|_p \,|\, \mathcal{L}(V) = \nu, \mathcal{L}(W) = \varrho\}, \quad \nu, \varrho \in \mathcal{M}_p^{\mathbb{R}},$$

where $\|V - W\|_p := (\mathbb{E}[|V - W|^p])^{1/p}$ is the usual L_p-norm. The spaces $(\mathcal{M}_p, \ell_p)$ and $(\mathcal{M}_p(\mu), \ell_p)$ for $1 \leq p < \infty$ are complete metric spaces, see [5]. The infimum in the definition of ℓ_p is a minimum. Random variables V', W' with distributions ν and ϱ respectively such that $\ell_p(\nu, \varrho) = \|V' - W'\|_p$ are called *optimal couplings*. They do exist for all $\nu, \varrho \in \mathcal{M}_1^{\mathbb{R}}$. We use the notation $\ell_p(X, Y) := \ell_p(\mathcal{L}(X), \mathcal{L}(Y))$ for random variables X and Y. Subsequently also the following inequality between the ℓ_p and ζ_s metrics is used:

$$\zeta_s(X, Y) \leq \Big((\mathbb{E}\,[|X|^s])^{1-1/s} + (\mathbb{E}\,[|Y|^s])^{1-1/s} \Big)\, \ell_s(X, Y), \quad 1 < s \leq 3, \tag{18}$$

where, for $1 < s \leq 2$, we need $\mathcal{L}(X), \mathcal{L}(Y) \in \mathcal{M}_s^{\mathbb{R}}(\mu)$ for some $\mu \in \mathbb{R}$ and, for $2 < s \leq 3$, we need $\mathcal{L}(X), \mathcal{L}(Y) \in \mathcal{M}_s^{\mathbb{R}}(\mu, \sigma^2)$ for some $\mu \in \mathbb{R}$ and $\sigma \geq 0$, see [8, Lemma 5.7].

On the product space $(\mathcal{M}_2^{\mathbb{R}}(0))^{\times d}$ we define

$$\ell_2^{\vee}((\nu_1, \dots, \nu_d), (\varrho_1, \dots, \varrho_d)) := \max_{1 \leq j \leq d} \ell_2(\nu_j, \varrho_j),$$

where $(\nu_1, \dots, \nu_d), (\mu_1, \dots, \mu_d) \in (\mathcal{M}_2^{\mathbb{R}}(0))^{\times d}$. Note that $(\mathcal{M}_2^{\mathbb{R}}(0))^{\times d}, \ell_2^{\vee})$ is a complete metric space as well.

Second, on the complex plane the minimal L_p-metric ℓ_p is defined similarly by

$$\ell_p(\nu, \varrho) := \inf\{\|V - W\|_p \,|\, \mathcal{L}(V) = \nu, \mathcal{L}(W) = \varrho\}, \quad \nu, \varrho \in \mathcal{M}_p^{\mathbb{C}},$$

with the analogous definition of the L_p-norm. The respective metric spaces are complete as in the real case and optimal couplings exist as well. On the product space $(\mathcal{M}_2^{\mathbb{C}}(0))^{\times d}$ we use

$$\ell_2^{\vee}((\nu_1, \dots, \nu_d), (\varrho_1, \dots, \varrho_d)) := \max_{1 \leq j \leq d} \ell_2(\nu_j, \varrho_j),$$

where $(\nu_1, \dots, \nu_d), (\mu_1, \dots, \mu_d) \in (\mathcal{M}_2^{\mathbb{C}}(0))^{\times d}$. Note that $(\mathcal{M}_2^{\mathbb{C}}(0))^{\times d}, \ell_2^{\vee})$ is a complete metric space as well.

Preview on the use of spaces and metrics. The guidance on which space and metric to use in which asymptotic regime of Pólya urns is as follows. We come back to the three types (a)–(c) of urns from the previous section:

(a) Urns that after scaling lead to convergence to a non-normal limit distribution. Typically such a convergence holds almost surely, however we only discuss convergence in distribution.

(b) Urns that after scaling lead to convergence to a normal limit. Such a convergence typically does not hold almost surely, but at least in distribution.

(c) Urns that eben after a proper scaling do not lead to convergence. Instead there is an asymptotic oscillatory behavior of the distributions. Such oscillatory behavior can even be captured almost surely, we discuss a (weak) description for distributions.

The cases of **type (a)** can be dealt with on the space $(\mathcal{M}_2^{\mathbb{R}}(\mu))^{\times d}$ with appropriate $\mu \in \mathbb{R}$ and $d \in \mathbb{N}$, where, by centering, one can always achieve the choice $\mu = 0$. One can either use the metric $\zeta_2^\vee$ or $\ell_2^\vee$ which lead to similar results, although based on different details in the proofs. We will only present the use of $\zeta_2^\vee$, since we do not see any advantage of $\ell_2^\vee$ here.

The cases of **type (b)** can be dealt with on the space $(\mathcal{M}_s^{\mathbb{R}}(\mu, \sigma^2))^{\times d}$ with $2 < s \leq 3$ and appropriate $\mu \in \mathbb{R}$, $\sigma > 0$ and $d \in \mathbb{N}$. By normalization, one can always achieve the choices $\mu = 0$ and $\sigma = 1$. Since in the context of urns third absolute moments in type (b) cases typically do exist, one can use $s = 3$ and the metric $\zeta_3^\vee$. We do not know how to use the $\ell_p^\vee$ metrics in type (b) cases.

The cases of **type (c)** can be dealt with on the space $(\mathcal{M}_2^{\mathbb{C}}(\gamma))^{\times d}$ with appropriate $\gamma \in \mathbb{R}$ and $d \in \mathbb{N}$. The metric of choice for the type (c) cases is the complex version of $\ell_2^\vee$. In our example below we will however use $\mathcal{M}_2^{\mathbb{C}}(\gamma_1) \times \cdots \times \mathcal{M}_2^{\mathbb{C}}(\gamma_d)$ with $\gamma_1, \ldots, \gamma_d \in \mathbb{C}$ to be able to work with a more natural scaling of the random variables, the metric still being $\ell_2^\vee$.

5 Associated fixed point equations

We fix $d, d' \in \mathbb{N}$, a $d \times d'$ matrix (A_{ir}) of random variables and a vector $(b_1, \ldots, b_d)$ of random variables. Either all of these random variables are real or all of them are complex. Furthermore, we are given a $d \times d'$ matrix $(\pi(ir))$ with all entries $\pi(ir) \in \{1, \ldots, d\}$. First, we consider the case, where all A_{ir} and all b_i are real. We associate a map

$$\begin{aligned} T : \left(\mathcal{M}^{\mathbb{R}}\right)^{\times d} &\to \left(\mathcal{M}^{\mathbb{R}}\right)^{\times d} \\ (\mu_1, \ldots, \mu_d) &\mapsto (T_1(\mu_1, \ldots, \mu_d), \ldots, T_d(\mu_1, \ldots, \mu_d)) \qquad (19) \\ T_i(\mu_1, \ldots, \mu_d) &:= \mathcal{L}\left(\sum_{r=1}^{d'} A_{ir} Z_{ir} + b_i\right) \qquad (20) \end{aligned}$$

with $(A_{i1}, \ldots, A_{id'}, b_i)$, $Z_{i1}, \ldots, Z_{id'}$ independent, and Z_{ir} distributed as $\mu_{\pi(ir)}$ for $r = 1, \ldots, d'$ and all components $i = 1, \ldots, d$.

In the case, where the A_{ir} and b_i are complex random variables, we define a map T' similar to T:

$$\begin{aligned} T' : \left(\mathcal{M}^{\mathbb{C}}\right)^{\times d} &\to \left(\mathcal{M}^{\mathbb{C}}\right)^{\times d} \qquad (21) \\ (\mu_1, \ldots, \mu_d) &\mapsto (T'_1(\mu_1, \ldots, \mu_d), \ldots, T'_d(\mu_1, \ldots, \mu_d)) \end{aligned}$$

with $T'_i(\mu_1, \ldots, \mu_d)$ defined as for T_i in (20).

For the three regimes discussed in the preview within section 4 we use the following three theorems (Theorem 5.1 for type (a), Theorem 5.2 for type (b), and Theorem 5.3 for type (c)) on existence of fixed-points of T and T'.

Theorem 5.1. *Assume that in the definition of* T *in (19) and (20) the* A_{ir} *and* b_i *are square integrable real random variables with* $\mathbb{E}[b_i] = 0$ *for all* $1 \leq i \leq d$ *and* $1 \leq r \leq d'$ *and*

$$\max_{1 \leq i \leq d} \sum_{r=1}^{d'} \mathbb{E}\left[A_{ir}^2\right] < 1. \tag{22}$$

Then the restriction of T *to* $(\mathcal{M}_2^{\mathbb{R}}(0))^{\times d}$ *has a unique fixed-point.*

Theorem 5.2. *Assume that in the definition of* T *in (19) and (20) for some* $\varepsilon > 0$ *the* A_{ir} *are* $L_{2+\varepsilon}$*-integrable real random variables and* $b_i = 0$ *for all* $1 \leq i \leq d$ *and* $1 \leq r \leq d'$*, that*

$$\sum_{r=1}^{d'} A_{ir}^2 = 1 \quad \text{for all } i = 1, \ldots, d \tag{23}$$

and

$$\min_{1 \leq i \leq d} \mathbb{P}\left(\max_{1 \leq r \leq d'} |A_{ir}| < 1\right) > 0. \tag{24}$$

Then, for all $\sigma^2 \geq 0$*, the restriction of* T *to* $(\mathcal{M}_{2+\varepsilon}^{\mathbb{R}}(0, \sigma^2))^{\times d}$ *has the unique fixed-point* $(\mathcal{N}(0, \sigma^2), \ldots, \mathcal{N}(0, \sigma^2))$.

Theorem 5.3. *Assume that in the definition of* T' *in (21) the* A_{ir} *and* b_i *are square integrable complex random variables for all* $1 \leq i \leq d$ *and* $1 \leq r \leq d'$ *and that for* $\gamma_1, \ldots, \gamma_d \in \mathbb{C}$ *we have*

$$\mathbb{E}[b_i] + \sum_{r=1}^{d'} \gamma_{\pi(ir)} \mathbb{E}[A_{ir}] = \gamma_i, \quad i = 1, \ldots, d. \tag{25}$$

If moreover

$$\max_{1 \leq i \leq d} \sum_{r=1}^{d'} \mathbb{E}\left[|A_{ir}|^2\right] < 1 \tag{26}$$

then the restriction of T' *to* $\mathcal{M}_2^{\mathbb{C}}(\gamma_1) \times \cdots \times \mathcal{M}_2^{\mathbb{C}}(\gamma_d)$ *has a unique fixed-point.*

Note that a special case of Theorem 5.1 was used in the proof of [12, Theorem 3.9 (iii)] with a similar proof technique as in our proof of Theorem 5.3.

The rest of this section contains the proofs of Theorems 5.1 – 5.3.

Proof. (Theorem 5.1). First note that for $(\mu_1, \ldots, \mu_d) \in (\mathcal{M}_2^{\mathbb{R}}(0))^{\times d}$, by independence in definition (20) and $\mathbb{E}[b_i] = 0$ we have $T_i(\mu_1, \ldots, \mu_d) \in \mathcal{M}_2^{\mathbb{R}}(0)$ for $i = 1, \ldots, d$. Hence, the restriction of T to $(\mathcal{M}_2^{\mathbb{R}}(0))^{\times d}$ maps into $(\mathcal{M}_2^{\mathbb{R}}(0))^{\times d}$.

Next, we show that the restriction of T to $(\mathcal{M}_2^{\mathbb{R}}(0))^{\times d}$ is a (strict) contraction with respect to the metric $\zeta_2^{\vee}$: For $(\mu_1, \ldots, \mu_d), (\nu_1, \ldots, \nu_d) \in (\mathcal{M}_2^{\mathbb{R}}(0))^{\times d}$ we first fix $i \in \{1, \ldots, d\}$. Let $Z_{i1}, \ldots, Z_{id'}$ and $Z'_{i1}, \ldots, Z'_{id'}$ be real random variables such that Z_{ir} is distributed as $\mu_{\pi(ir)}$ and Z'_{ir} is distributed as $\nu_{\pi(ir)}$. Moreover, assume that both families $\{(A_{i1}, \ldots, A_{id'}, b_i), Z_{i1}, \ldots, Z_{id'}\}$ and $\{(A_{i1}, \ldots, A_{id'}, b_i), Z'_{i1}, \ldots, Z'_{id'}\}$ are independent. Then we have

$$T_i(\mu_1, \ldots, \mu_d) = \mathcal{L}\left(\sum_{r=1}^{d'} A_{ir} Z_{ir} + b_i\right), \quad T_i(\nu_1, \ldots, \nu_d) = \mathcal{L}\left(\sum_{r=1}^{d'} A_{ir} Z'_{ir} + b_i\right). \tag{27}$$

Conditioning on $(A_{i1},\dots,A_{id'},b_i)$ and denoting this vector's distribution by Υ we obtain

$$\begin{aligned}
&\zeta_2(T_i(\mu_1,\dots,\mu_d),T_i(\nu_1,\dots,\nu_d))\\
&\quad=\sup_{f\in\mathcal{F}_2}\left|\int f\left(\sum_{r=1}^{d'}\alpha_r Z_{ir}+\beta\right)-f\left(\sum_{r=1}^{d'}\alpha_r Z'_{ir}+\beta\right)d\Upsilon(\alpha_1,\dots,\alpha_{d'},\beta)\right|\\
&\quad\le\int\sup_{f\in\mathcal{F}_2}\left|f\left(\sum_{r=1}^{d'}\alpha_r Z_{ir}+\beta\right)-f\left(\sum_{r=1}^{d'}\alpha_r Z'_{ir}+\beta\right)\right|d\Upsilon(\alpha_1,\dots,\alpha_{d'},\beta)\\
&\quad=\int\zeta_2\left(\sum_{r=1}^{d'}\alpha_r Z_{ir}+\beta,\sum_{r=1}^{d'}\alpha_r Z'_{ir}+\beta\right)d\Upsilon(\alpha_1,\dots,\alpha_{d'},\beta) \qquad (28)
\end{aligned}$$

Since ζ_2 is $(2,+)$-ideal, we obtain from (15) that $\zeta_2(\sum\alpha_r Z_{ir}+\beta,\sum\alpha_r Z'_{ir}+\beta)\le\sum\alpha_r^2\zeta_2(Z_{ir},Z'_{ir})$. Hence, we can further estimate

$$\begin{aligned}
&\zeta_2(T_i(\mu_1,\dots,\mu_d),T_i(\nu_1,\dots,\nu_d))\\
&\quad\le\int\sum_{r=1}^{d'}\alpha_r^2\zeta_2(Z_{ir},Z'_{ir})\,d\Upsilon(\alpha_1,\dots,\alpha_{d'},\beta)\\
&\quad=\int\sum_{r=1}^{d'}\alpha_r^2\zeta_2(\mu_{\pi(ir)},\nu_{\pi(ir)})\,d\Upsilon(\alpha_1,\dots,\alpha_{d'},\beta)\\
&\quad\le\left(\sum_{r=1}^{d'}\mathbb{E}\left[A_{ir}^2\right]\right)\zeta_2^\vee((\mu_1,\dots,\mu_d),(\nu_1,\dots,\nu_d)). \qquad (29)
\end{aligned}$$

Now, taking the maximum over i yields

$$\zeta_2^\vee(T(\mu_1,\dots,\mu_d),T(\nu_1,\dots,\nu_d))\le\left(\max_{1\le i\le d}\sum_{r=1}^{d'}\mathbb{E}\left[A_{ir}^2\right]\right)\zeta_2^\vee((\mu_1,\dots,\mu_d),(\nu_1,\dots,\nu_d)). \qquad (30)$$

Hence, condition (22) implies that the restriction of T to $(\mathcal{M}_2^{\mathbb{R}}(0))^{\times d}$ is a contraction. Since the metric $\zeta_2^\vee$ is complete, Banach's fixed-point theorem implies the assertion. □

Proof. (Theorem 5.2). This proof is similar to the previous proof of Theorem 5.1. Let $\varepsilon>0$ be as in Theorem 5.2 and $\sigma>0$ be arbitrary. First note that for $(\mu_1,\dots,\mu_d)\in(\mathcal{M}_{2+\varepsilon}^{\mathbb{R}}(0,\sigma^2))^{\times d}$, by independence in definition (20), condition (23), and $b_i=0$ we have $T_i(\mu_1,\dots,\mu_d)\in\mathcal{M}_{2+\varepsilon}^{\mathbb{R}}(0,\sigma^2)$ for $i=1,\dots,d$. Hence, the restriction of T to $(\mathcal{M}_{2+\varepsilon}^{\mathbb{R}}(0,\sigma^2))^{\times d}$ maps into $(\mathcal{M}_{2+\varepsilon}^{\mathbb{R}}(0,\sigma^2))^{\times d}$.

We set $s:=(2+\varepsilon)\wedge 3$. For $(\mu_1,\dots,\mu_d),(\nu_1,\dots,\nu_d)\in(\mathcal{M}_{2+\varepsilon}^{\mathbb{R}}(0,\sigma^2))^{\times d}$ we choose $Z_{i1},\dots,Z_{id'}$ and $Z'_{i1},\dots,Z'_{id'}$ as in the proof of Theorem 5.1 such that we have (27). Note that with our choice of s we have $\zeta_s(T_i(\mu_1,\dots,\mu_d),T_i(\nu_1,\dots,\nu_d))<\infty$. With an estimate analogous to (28) – (30), using now that ζ_s is $(s,+)$-ideal, we obtain

$$\zeta_s^\vee(T(\mu_1,\dots,\mu_d),T(\nu_1,\dots,\nu_d))\le\left(\max_{1\le i\le d}\sum_{r=1}^{d'}\mathbb{E}\left[|A_{ir}|^s\right]\right)\zeta_s^\vee((\mu_1,\dots,\mu_d),(\nu_1,\dots,\nu_d)).$$

Note that $s > 2$ and the conditions (23) and (24) imply that $\sum_{r=1}^{d'} \mathbb{E}[|A_{ir}|^s] < 1$ for all $i = 1, \dots, d$. Hence, the restriction of T to $(\mathcal{M}_{2+\varepsilon}^{\mathbb{R}}(0, \sigma^2))^{\times d}$ is a contraction and the completeness of $\zeta_s^\vee$ implies the existence of a unique fixed-point. With the convolution property $\mathcal{N}(0, \sigma_1^2) * \mathcal{N}(0, \sigma_2^2) = \mathcal{N}(0, \sigma_1^2 + \sigma_2^2)$ for $\sigma_1, \sigma_2 \geq 0$ one can directly check that $(\mathcal{N}(0, \sigma^2), \dots, \mathcal{N}(0, \sigma^2))$ is a fixed-point of T in $(\mathcal{M}_{2+\varepsilon}^{\mathbb{R}}(0, \sigma^2))^{\times d}$. □

Proof. (Theorem 5.3). Let $\gamma_1, \dots, \gamma_d$ be as in Theorem 5.3 and abbreviate $\mathcal{P} := \mathcal{M}_2^{\mathbb{C}}(\gamma_1) \times \cdots \times \mathcal{M}_2^{\mathbb{C}}(\gamma_d)$. First note that for $(\mu_1, \dots, \mu_d) \in \mathcal{P}$ from independence in the definition of $T_i'(\mu_1, \dots, \mu_d)$ and the finite second moments of the A_{ir} and b_i we obtain $T_i'(\mu_1, \dots, \mu_d) \in \mathcal{M}_2^{\mathbb{C}}$ for all $i = 1, \dots, d$. For a random variable W with distribution $T_i'(\mu_1, \dots, \mu_d)$ we have

$$\mathbb{E}[W] = \sum_{r=1}^{d'} \mathbb{E}[A_{ir}]\gamma_{\pi(ir)} + \mathbb{E}[b_i] = \gamma_i$$

by condition (25). Hence, the restriction of T' to $\mathcal{P}$ maps into $\mathcal{P}$.

Next, we show that the restriction of T' to $\mathcal{P}$ is a contraction with respect to the metric $\ell_2^\vee$: For $(\mu_1, \dots, \mu_d), (\nu_1, \dots, \nu_d) \in \mathcal{P}$ we first fix $i \in \{1, \dots, d\}$. Let (Z_{ir}, Z_{ir}') be an optimal coupling of $\mu_{\pi(ir)}$ and $\nu_{\pi(ir)}$ for $r = 1, \dots, d'$ such that $(Z_{i1}, Z_{i1}'), \dots, (Z_{id'}, Z_{id'}'), (A_{i1}, \dots, A_{id'}, b_i)$ are independent. Then we have

$$T_i'(\mu_1, \dots, \mu_d) = \mathcal{L}\left(\sum_{r=1}^{d'} A_{ir} Z_{ir} + b_i\right), \quad T_i'(\nu_1, \dots, \nu_d) = \mathcal{L}\left(\sum_{r=1}^{d'} A_{ir} Z_{ir}' + b_i\right). \tag{31}$$

Denoting by $\overline{\gamma}$ the complex conjugate of $\gamma \in \mathbb{C}$ we obtain

$$\begin{aligned}
&\ell_2^2(T_i'(\mu_1, \dots, \mu_d), T_i'(\nu_1, \dots, \nu_d)) \\
&\leq \mathbb{E}\left[\left|\sum_{r=1}^{d'} A_{ir}(Z_{ir} - Z_{ir}')\right|^2\right] \\
&= \mathbb{E}\left[\sum_{r=1}^{d'} |A_{ir}|^2 |Z_{ir} - Z_{ir}'|^2\right] + \mathbb{E}\left[\sum_{r \neq t} A_{ir}(Z_{ir} - Z_{ir}')\overline{A_{it}(Z_{it} - Z_{it}')}\right] \\
&= \sum_{r=1}^{d'} \mathbb{E}\left[|A_{ir}|^2\right] \ell_2^2(\mu_{\pi(ir)}, \nu_{\pi(ir)}) \qquad (32) \\
&\leq \left(\sum_{r=1}^{d'} \mathbb{E}\left[|A_{ir}|^2\right]\right) \left(\ell_2^\vee((\mu_1, \dots, \mu_d), (\nu_1, \dots, \nu_d))\right)^2.
\end{aligned}$$

For equality (32) we firstly use that $Z_{ir} - Z_{ir}'$ and $Z_{it} - Z_{it}'$ are independent, centered factors, so that the expectation of the sum over $r \neq t$ is 0 and secondly that (Z_{ir}, Z_{ir}') are optimal couplings of $(\mu_{\pi(ir)}, \nu_{\pi(ir)})$ such that $\mathbb{E}[|Z_{ir} - Z_{ir}'|^2] = \ell_2^2(\mu_{\pi(ir)}, \nu_{\pi(ir)})$.

Now, taking the maximum over i yields

$$\ell_2^\vee(T'(\mu_1, \dots, \mu_d), T'(\nu_1, \dots, \nu_d)) \leq \left(\max_{1 \leq i \leq d} \sum_{r=1}^{d'} \mathbb{E}\left[|A_{ir}|^2\right]\right)^{1/2} \ell_2^\vee((\mu_1, \dots, \mu_d), (\nu_1, \dots, \nu_d)).$$

Hence, condition (26) implies that the restriction of T' to $\mathcal{P}$ is a contraction. Since the metric $\ell_2^\vee$ is complete, Banach's fixed-point theorem implies the assertion. □

6 Convergence and examples

In this section a couple of concrete Pólya urns are considered and convergence of the normalized numbers of balls of a color is shown within the product metrics defined in section 4. The proofs are generic such that they can easily be transferred to other urns of the types (a)–(c) in section 3. We always show limit laws for the initial compositions of the urn with one ball of (arbitrary) color. Limit laws for other initial compositions can be obtained from these by appropriate convolution with coefficients which are powers of components of an independent Dirichlet distributed vector. We leave the details to the reader.

6.1 2 × 2 deterministic replacement urns

A discussion of urns with a general balanced 2×2 replacement matrix as in (1) is given in Bagchi and Pal [2]. Subsequently, we assume the conditions in (1) and, as in [2], that $bc > 0$. As shown in [2] asymptotic normal behavior occurs for these urns when $a - c \leq (a+b)/2$ (type (b) in section 4), whereas $a - c > (a+b)/2$ leads to limit laws with non-normal limit distributions (type (a) in section 4). In this section we show how to derive these results by our contraction approach. With B_n^{b} and B_n^{w} as in the beginning of section 2 we denote expectations by $\mu_{\mathrm{b}}(n)$ and $\mu_{\mathrm{w}}(n)$. These values can be derived exactly, see [2],

$$\mu_{\mathrm{b}}(n) = \frac{c(a+b)}{b+c} n + \frac{b\,\Gamma\left(\frac{1}{a+b}\right)}{(b+c)\Gamma\left(\frac{1+a-c}{a+b}\right)} \frac{\Gamma\left(n + \frac{1+a-c}{a+b}\right)}{\Gamma\left(n + \frac{1}{a+b}\right)} + \frac{c}{b+c}, \tag{33}$$

$$\mu_{\mathrm{w}}(n) = \frac{c(a+b)}{b+c} n - \frac{c\,\Gamma\left(\frac{1}{a+b}\right)}{(b+c)\Gamma\left(\frac{1+a-c}{a+b}\right)} \frac{\Gamma\left(n + \frac{1+a-c}{a+b}\right)}{\Gamma\left(n + \frac{1}{a+b}\right)} + \frac{c}{b+c}. \tag{34}$$

Non-normal limit case. We first discuss the non-normal case $a - c > (a+b)/2$. Note that with $\lambda := (a-c)/(a+b)$ and excluding the case $bc = 0$, we have $1/2 < \lambda < 1$ and, as $n \to \infty$,

$$\mu_{\mathrm{b}}(n) = c_{\mathrm{b}} n + d_{\mathrm{b}} n^{\lambda} + o(n^{\lambda}), \quad \mu_{\mathrm{w}}(n) = c_{\mathrm{w}} n + d_{\mathrm{w}} n^{\lambda} + o(n^{\lambda}) \tag{35}$$

with

$$c_{\mathrm{b}} = c_{\mathrm{w}} = \frac{c(a+b)}{b+c}, \quad d_{\mathrm{b}} = \frac{b\,\Gamma\left(\frac{1}{a+b}\right)}{(b+c)\Gamma\left(\frac{1+a-c}{a+b}\right)}, \quad d_{\mathrm{w}} = -\frac{c\,\Gamma\left(\frac{1}{a+b}\right)}{(b+c)\Gamma\left(\frac{1+a-c}{a+b}\right)}. \tag{36}$$

We use the normalizations $X_0 := Y_0 := 0$ and

$$X_n := \frac{B_n^{\mathrm{b}} - \mu_{\mathrm{b}}(n)}{n^{\lambda}}, \quad Y_n := \frac{B_n^{\mathrm{w}} - \mu_{\mathrm{w}}(n)}{n^{\lambda}}, \quad n \geq 1. \tag{37}$$

Note that we do not have to identify the order of the variance in advance. It turns out that it is sufficient to use the order of the error terms $d_{\mathrm{b}} n^{\lambda}$ and $d_{\mathrm{w}} n^{\lambda}$ in the expansions (35). From the system (2)–(3) we obtain for the scaled quantities X_n, Y_n the system, for $n \geq 1$,

$$X_n \stackrel{d}{=} \sum_{r=1}^{a+1} \left(\frac{I_r^{(n)}}{n}\right)^{\lambda} X_{I_r^{(n)}}^{(r)} + \sum_{r=a+2}^{K} \left(\frac{I_r^{(n)}}{n}\right)^{\lambda} Y_{I_r^{(n)}}^{(r)} + b_{\mathrm{b}}(n), \tag{38}$$

$$Y_n \stackrel{d}{=} \sum_{r=1}^{c} \left(\frac{I_r^{(n)}}{n}\right)^{\lambda} X_{I_r^{(n)}}^{(r)} + \sum_{r=c+1}^{K} \left(\frac{I_r^{(n)}}{n}\right)^{\lambda} Y_{I_r^{(n)}}^{(r)} + b_{\mathrm{w}}(n), \tag{39}$$

with

$$b_{\mathrm{b}}(n) = d_{\mathrm{b}} \left(-1 + \sum_{r=1}^{a+1} \left(\frac{I_r^{(n)}}{n} \right)^{\lambda} \right) + d_{\mathrm{w}} \sum_{r=a+2}^{K} \left(\frac{I_r^{(n)}}{n} \right)^{\lambda} + o(1), \tag{40}$$

$$b_{\mathrm{w}}(n) = d_{\mathrm{b}} \sum_{r=1}^{c} \left(\frac{I_r^{(n)}}{n} \right)^{\lambda} + d_{\mathrm{w}} \left(-1 + \sum_{r=c+1}^{K} \left(\frac{I_r^{(n)}}{n} \right)^{\lambda} \right) + o(1), \tag{41}$$

with conditions on independence and identical distributions analogously to (2) and (3). In view of Lemma 2.1 this suggests for limits X and Y of X_n and Y_n respectively

$$X \stackrel{d}{=} \sum_{r=1}^{a+1} D_r^{\lambda} X^{(r)} + \sum_{r=a+2}^{K} D_r^{\lambda} Y^{(r)} + b_{\mathrm{b}}, \tag{42}$$

$$Y \stackrel{d}{=} \sum_{r=1}^{c} D_r^{\lambda} X^{(r)} + \sum_{r=c+1}^{K} D_r^{\lambda} Y^{(r)} + b_{\mathrm{w}}, \tag{43}$$

with

$$b_{\mathrm{b}} = d_{\mathrm{b}} \left(-1 + \sum_{r=1}^{a+1} D_r^{\lambda} \right) + d_{\mathrm{w}} \sum_{r=a+2}^{K} D_r^{\lambda},$$

$$b_{\mathrm{w}} = d_{\mathrm{b}} \sum_{r=1}^{c} D_r^{\lambda} + d_{\mathrm{w}} \left(-1 + \sum_{r=c+1}^{K} D_r^{\lambda} \right),$$

where $(D_1, \ldots, D_K)$, $X^{(1)}, \ldots, X^{(K)}$, $Y^{(1)}, \ldots, Y^{(K)}$ are independent, and the $X^{(r)}$ are distributed as X, the $Y^{(r)}$ are distributed as Y and $(D_1, \ldots, D_K)$ is as in Lemma 2.1. Note that the moments $\mathbb{E}[D_r^{\lambda}]$ and the form of d_{b} and d_{w} in (36) imply $\mathbb{E}[b_{\mathrm{b}}] = \mathbb{E}[b_{\mathrm{w}}] = 0$. From $\lambda > 1/2$ and $\sum_{r=1}^{K} D_r = 1$ we obtain

$$\sum_{r=1}^{K} \mathbb{E}\left[D_r^{2\lambda} \right] < 1.$$

Hence, Theorem 5.1 applies to the map associated to the system (42)–(43) and implies that there exists a unique solution $(\mathcal{L}(\Lambda_{\mathrm{b}}), \mathcal{L}(\Lambda_{\mathrm{w}}))$ in the space $\mathcal{M}_2^{\mathbb{R}}(0) \times \mathcal{M}_2^{\mathbb{R}}(0)$ to (42)–(43). The following convergence proof resembles ideas from Neininger and Rüschendorf [20].

Theorem 6.1. *Consider the Pólya urn with replacement matrix (1) with* $a - c > (a+b)/2$ *and* $bc > 0$ *and the normalized numbers* X_n *and* Y_n *of black balls as in (37). Furthermore let* $(\mathcal{L}(\Lambda_{\mathrm{b}}), \mathcal{L}(\Lambda_{\mathrm{w}}))$ *denote the in* $\mathcal{M}_2^{\mathbb{R}}(0) \times \mathcal{M}_2^{\mathbb{R}}(0)$ *unique solution of (42)–(43). Then, as* $n \to \infty$,

$$\zeta_2^{\vee}\left((X_n, Y_n), (\Lambda_{\mathrm{b}}, \Lambda_{\mathrm{w}}) \right) \to 0.$$

In particular, as $n \to \infty$,

$$X_n \xrightarrow{d} \Lambda_{\mathrm{b}}, \quad Y_n \xrightarrow{d} \Lambda_{\mathrm{w}}. \tag{44}$$

Proof. We first define, for $n \geq 1$, the accompanying sequences

$$Q_n^{\mathrm{b}} := \sum_{r=1}^{a+1} \left(\frac{I_r^{(n)}}{n} \right)^{\lambda} \Lambda_{\mathrm{b}}^{(r)} + \sum_{r=a+2}^{K} \left(\frac{I_r^{(n)}}{n} \right)^{\lambda} \Lambda_{\mathrm{w}}^{(r)} + b_{\mathrm{b}}(n), \tag{45}$$

$$Q_n^{\mathrm{w}} := \sum_{r=1}^{c} \left(\frac{I_r^{(n)}}{n} \right)^{\lambda} \Lambda_{\mathrm{b}}^{(r)} + \sum_{r=c+1}^{K} \left(\frac{I_r^{(n)}}{n} \right)^{\lambda} \Lambda_{\mathrm{w}}^{(r)} + b_{\mathrm{w}}(n), \tag{46}$$

with $b_{\mathrm{b}}(n)$ and $b_{\mathrm{w}}(n)$ as in (40) and the $\Lambda_{\mathrm{b}}^{(r)}$, $\Lambda_{\mathrm{b}}^{(r)}$ and $I^{(n)}$ being independent, where the $\Lambda_{\mathrm{b}}^{(r)}$ are distributed as Λ_{b} and the $\Lambda_{\mathrm{w}}^{(r)}$ are distributed as Λ_{w} for the respective values of r. Note that Q_n^{b} and Q_n^{w} are centered with finite second moment since $\mathcal{L}(\Lambda_{\mathrm{b}}), \mathcal{L}(\Lambda_{\mathrm{b}}) \in \mathcal{M}_2^{\mathbb{R}}(0)$. Hence, ζ_2 distances between $X_n, Y_n, Q_n^{\mathrm{b}}, Q_n^{\mathrm{w}}, \Lambda_{\mathrm{b}}$ and Λ_{w} are finite. To bound

$$\Delta(n) := \zeta_2^{\vee}((X_n, Y_n), (\Lambda_{\mathrm{b}}, \Lambda_{\mathrm{w}}))$$

we look at the distances

$$\Delta_{\mathrm{b}}(n) := \zeta_2(X_n, \Lambda_{\mathrm{b}}), \quad \Delta_{\mathrm{w}}(n) := \zeta_2(Y_n, \Lambda_{\mathrm{w}}).$$

We start with the estimate

$$\zeta_2(X_n, \Lambda_{\mathrm{b}}) \leq \zeta_2(X_n, Q_n^{\mathrm{b}}) + \zeta_2(Q_n^{\mathrm{b}}, \Lambda_{\mathrm{b}}). \tag{47}$$

We first show for the second summand in the latter display that $\zeta_2(Q_n^{\mathrm{b}}, \Lambda_{\mathrm{b}}) \to 0$ as $n \to \infty$: With inequality (18) we have

$$\zeta_2(Q_n^{\mathrm{b}}, \Lambda_{\mathrm{b}}) \leq (\|Q_n^{\mathrm{b}}\|_2 + \|\Lambda_{\mathrm{b}}\|_2) \ell_2(Q_n^{\mathrm{b}}, \Lambda_{\mathrm{b}}).$$

Moreover, $\|\Lambda_{\mathrm{b}}\|_2 < \infty$ since $\mathcal{L}(\Lambda_{\mathrm{b}}) \in \mathcal{M}_2^{\mathbb{R}}$ and, by definition of Q_n^{b} and with $|I_r^{(n)}/n| \leq 1$ we have that $\|Q_n^{\mathrm{b}}\|_2$ is uniformly bounded in n. Hence, it is sufficient to show $\ell_2(Q_n^{\mathrm{b}}, \Lambda_{\mathrm{b}}) \to 0$. We have, using the independence properties in (45) and (42), that

$$\begin{aligned} &\ell_2(Q_n^{\mathrm{b}}, \Lambda_{\mathrm{b}}) \\ &\leq \sum_{r=1}^{a+1} \left\| \left(\frac{I_r^{(n)}}{n} \right)^{\lambda} - D_r^{\lambda} \right\|_2 \|\Lambda_{\mathrm{b}}^{(r)}\|_2 + \sum_{r=a+2}^{K} \left\| \left(\frac{I_r^{(n)}}{n} \right)^{\lambda} - D_r^{\lambda} \right\|_2 \|\Lambda_{\mathrm{w}}^{(r)}\|_2 + \|b_{\mathrm{b}}(n) - b_{\mathrm{b}}\|_2. \end{aligned}$$

Lemma 2.1 implies that $\|(I_r^{(n)}/n)^{\lambda} - D_r^{\lambda}\|_2 \to 0$ as $n \to \infty$, which as well implies $\|b_{\mathrm{b}}(n) - b_{\mathrm{b}}\|_2 \to 0$. Hence, we obtain $\ell_2(Q_n^{\mathrm{b}}, \Lambda_{\mathrm{b}}) \to 0$ and $\zeta_2(Q_n^{\mathrm{b}}, \Lambda_{\mathrm{b}}) \to 0$.

Next, we bound the first summand $\zeta_2(X_n, Q_n^{\mathrm{b}})$ in (47). We condition on $I^{(n)}$. Note that conditionally on $I^{(n)}$ we have that $b_{\mathrm{b}}(n)$ is deterministic, which, for integration, we denote by

$\beta = \beta(I^{(n)})$. Denoting the distribution of $I^{(n)}$ by Υ_n and $\mathbf{i} := (i_1, \dots, i_K)$ this yields

$$\begin{aligned}
\zeta_2(X_n, Q_n^{\mathrm{b}}) \le \int \zeta_2 \Bigg(&\sum_{r=1}^{a+1} \left(\frac{i_r}{n}\right)^{\lambda} X_{i_r}^{(r)} + \sum_{r=a+2}^{K} \left(\frac{i_r}{n}\right)^{\lambda} Y_{i_r}^{(r)} + \beta, \\
&\sum_{r=1}^{a+1} \left(\frac{i_r}{n}\right)^{\lambda} \Lambda_{\mathrm{b}}^{(r)} + \sum_{r=a+2}^{K} \left(\frac{i_r}{n}\right)^{\lambda} \Lambda_{\mathrm{w}}^{(r)} + \beta \Bigg)\, d\Upsilon_n(\mathbf{i}) \\
\le \int &\sum_{r=1}^{a+1} \left(\frac{i_r}{n}\right)^{2\lambda} \zeta_2(X_{i_r}^{(r)}, \Lambda_{\mathrm{b}}^{(r)}) + \sum_{r=a+2}^{K} \left(\frac{i_r}{n}\right)^{2\lambda} \zeta_2(Y_{i_r}^{(r)}, \Lambda_{\mathrm{w}}^{(r)})\, d\Upsilon_n(\mathbf{i}) \qquad (48) \\
= &\sum_{r=1}^{a+1} \mathbb{E}\left[\left(\frac{I_r^{(n)}}{n}\right)^{2\lambda} \Delta_{\mathrm{b}}(I_r^{(n)})\right] + \sum_{r=a+2}^{K} \mathbb{E}\left[\left(\frac{I_r^{(n)}}{n}\right)^{2\lambda} \Delta_{\mathrm{w}}(I_r^{(n)})\right] \\
\le &\sum_{r=1}^{K} \mathbb{E}\left[\left(\frac{I_r^{(n)}}{n}\right)^{2\lambda} \Delta(I_r^{(n)})\right],
\end{aligned}$$

where, for (48) we use that ζ_2 is $(2,+)$ ideal, as well as (16). Altogether, the estimate started in (47) yields

$$\Delta_{\mathrm{b}}(n) \le \sum_{r=1}^{K} \mathbb{E}\left[\left(\frac{I_r^{(n)}}{n}\right)^{2\lambda} \Delta(I_r^{(n)})\right] + o(1).$$

With the same argument we obtain the same upper bound for $\Delta_{\mathrm{w}}(n)$. Thus, using also that $I_1^{(n)}, \dots, I_K^{(n)}$ are identically distributed we have

$$\Delta(n) \le K\mathbb{E}\left[\left(\frac{I_1^{(n)}}{n}\right)^{2\lambda} \Delta(I_1^{(n)})\right] + o(1). \qquad (49)$$

Now, a standard argument implies $\Delta(n) \to 0$ as follows: First from (49) we obtain with $I_1^{(n)}/n \to D_1$ in L_2 and, by $\lambda > 1/2$, with $\vartheta := K\mathbb{E}[D_1^{2\lambda}] < 1$ that

$$\begin{aligned}
\Delta(n) &\le K\mathbb{E}\left[\left(\frac{I_1^{(n)}}{n}\right)^{2\lambda}\right] \max_{0\le k\le n-1} \Delta(k) + o(1) \\
&\le (\vartheta + o(1)) \max_{0\le k\le n-1} \Delta(k) + o(1).
\end{aligned}$$

Since $\vartheta < 1$ this implies that the sequence $(\Delta(n))_{n\ge0}$ is bounded. We denote $\eta := \sup_{n\ge0} \Delta(n)$ and $\xi := \limsup_{n\to\infty} \Delta(n)$. For any $\varepsilon > 0$ there exists an $n_0 \ge 0$ such that $\Delta(n) \le \xi + \varepsilon$ for all $n \ge n_0$. Hence, from (49) we obtain

$$\Delta(n) \le K\mathbb{E}\left[\mathbf{1}_{\left\{I_1^{(n)} < n_0\right\}} \left(\frac{I_1^{(n)}}{n}\right)^{2\lambda}\right] \eta + K\mathbb{E}\left[\mathbf{1}_{\left\{I_1^{(n)} \ge n_0\right\}} \left(\frac{I_1^{(n)}}{n}\right)^{2\lambda}\right] (\xi + \varepsilon) + o(1).$$

With $n \to \infty$ this implies

$$\xi \le \vartheta(\xi + \varepsilon).$$

Since $\vartheta < 1$ and $\varepsilon > 0$ is arbitrary this implies $\xi = 0$. Hence, we have $\zeta_2^\vee\left((X_n, Y_n), (\Lambda_\mathrm{b}, \Lambda_\mathrm{w})\right) \to 0$ as $n \to \infty$. Since convergence in ζ_2 implies weak convergence, this implies (44) as well. $\square$

The normal limit case. Now, we discuss the normal limit case $a - c \le (a+b)/2$, where we first consider $a - c < (a+b)/2$. (The remaining case $a - c = (a+b)/2$ is similar with more involved expansions for the first two moments.) The formulae (33), (34) now imply

$$\mu_\mathrm{b}(n) = c_\mathrm{b} n + o(\sqrt{n}), \quad \mu_\mathrm{w}(n) = c_\mathrm{w} n + o(\sqrt{n}) \tag{50}$$

with c_b and c_w as in (36). As it is usual in the use of the contraction method for proving normal limit laws based on the metric ζ_3 we need also an expansion of the variance. We denote the variances of B_n^b and B_n^w by $\sigma_\mathrm{b}^2(n)$ and $\sigma_\mathrm{w}^2(n)$. Additionally to $bc = 0$ we exclude the case $a = c$. (In this case there is a trivial non-random evolution of the urn). From [2] we have as $n \to \infty$:

$$\sigma_\mathrm{b}^2(n) = f_\mathrm{b} n + o(n), \quad \sigma_\mathrm{w}^2(n) = f_\mathrm{w} n + o(n), \tag{51}$$

with

$$f_\mathrm{b} = f_\mathrm{w} = \frac{bc(a-c)^2}{(a + b - 2\frac{a-c}{a+b})(a+b)(b+c)} > 0.$$

We use the normalizations $X_0 := Y_0 := X_1 := Y_1 := 0$ and

$$X_n := \frac{B_n^\mathrm{b} - \mu_\mathrm{b}(n)}{\sigma_\mathrm{b}(n)}, \quad Y_n := \frac{B_n^\mathrm{w} - \mu_\mathrm{w}(n)}{\sigma_\mathrm{w}(n)}, \quad n \ge 2. \tag{52}$$

From the system (2)–(3) we obtain for the scaled quantities X_n, Y_n the system, for $n \ge 1$,

$$X_n \stackrel{d}{=} \sum_{r=1}^{a+1} \frac{\sigma_\mathrm{b}(I_r^{(n)})}{\sigma_\mathrm{b}(n)} X_{I_r^{(n)}}^{(r)} + \sum_{r=a+2}^{K} \frac{\sigma_\mathrm{w}(I_r^{(n)})}{\sigma_\mathrm{w}(n)} Y_{I_r^{(n)}}^{(r)} + e_\mathrm{b}(n), \tag{53}$$

$$Y_n \stackrel{d}{=} \sum_{r=1}^{c} \frac{\sigma_\mathrm{b}(I_r^{(n)})}{\sigma_\mathrm{b}(n)} X_{I_r^{(n)}}^{(r)} + \sum_{r=c+1}^{K} \frac{\sigma_\mathrm{w}(I_r^{(n)})}{\sigma_\mathrm{w}(n)} Y_{I_r^{(n)}}^{(r)} + e_\mathrm{w}(n), \tag{54}$$

with conditions on independence and identical distributions analogously to (2) and (3). We have $\|e_\mathrm{b}(n)\|_\infty, \|e_\mathrm{w}(n)\|_\infty \to 0$ since the leading linear terms in the expansions (50) cancel out and the error terms $o(\sqrt{n})$ are asymptotically eliminated by the scaling of order $1/\sqrt{n}$. In view of Lemma 2.1 this suggests for limits X and Y of X_n and Y_n respectively

$$X \stackrel{d}{=} \sum_{r=1}^{a+1} \sqrt{D_r} X^{(r)} + \sum_{r=a+2}^{K} \sqrt{D_r} Y^{(r)}, \tag{55}$$

$$Y \stackrel{d}{=} \sum_{r=1}^{c} \sqrt{D_r} X^{(r)} + \sum_{r=c+1}^{K} \sqrt{D_r} Y^{(r)}, \tag{56}$$

where $(D_1, \ldots, D_K)$, $X^{(1)}, \ldots, X^{(K)}$, $Y^{(1)}, \ldots, Y^{(K)}$ are independent, and the $X^{(r)}$ are distributed as X and the $Y^{(r)}$ are distributed as Y. To the map associated to the system (55)–(56) we can apply Theorem 5.2. The conditions (23) and (24) are trivially satisfied. Hence $(\mathcal{N}(0,1), \mathcal{N}(0,1))$ is the unique fixed-point of the associated map in the space $\mathcal{M}_3^\mathbb{R}(0,1) \times \mathcal{M}_3^\mathbb{R}(0,1)$.

Theorem 6.2. *Consider the Pólya urn with replacement matrix (1) with* $a - c < (a+b)/2$ *and* $bc > 0$ *and the normalized numbers* X_n *and* Y_n *of black balls as in (52). Then, as* $n \to \infty$*,*

$$\zeta_3^{\vee}\left((X_n, Y_n), (\mathcal{N}(0,1), \mathcal{N}(0,1))\right) \to 0.$$

In particular, as $n \to \infty$*,*

$$X_n \xrightarrow{d} \mathcal{N}(0,1), \quad Y_n \xrightarrow{d} \mathcal{N}(0,1). \tag{57}$$

Proof. The proof of this Theorem can be done along the lines of the proof of Theorem 6.1. However, more care has to be taken in the definition of the quantities corresponding to Q_n^{b} and Q_n^{w} in (45) in order to assure finiteness of the ζ_3 distances. A possible choice is, for $n \geq 2$,

$$\tilde{Q}_n^{\mathrm{b}} := \sum_{r=1}^{a+1} \mathbf{1}_{\left\{I_r^{(n)} \geq 2\right\}} \frac{\sigma_{\mathrm{b}}(I_r^{(n)})}{\sigma_{\mathrm{b}}(n)} N_r + \sum_{r=a+2}^{K} \mathbf{1}_{\left\{I_r^{(n)} \geq 2\right\}} \frac{\sigma_{\mathrm{w}}(I_r^{(n)})}{\sigma_{\mathrm{w}}(n)} N_r + e_{\mathrm{b}}(n), \tag{58}$$

$$\tilde{Q}_n^{\mathrm{w}} \stackrel{d}{=} \sum_{r=1}^{c} \mathbf{1}_{\left\{I_r^{(n)} \geq 2\right\}} \frac{\sigma_{\mathrm{b}}(I_r^{(n)})}{\sigma_{\mathrm{b}}(n)} N_r + \sum_{r=c+1}^{K} \mathbf{1}_{\left\{I_r^{(n)} \geq 2\right\}} \frac{\sigma_{\mathrm{w}}(I_r^{(n)})}{\sigma_{\mathrm{w}}(n)} N_r + e_{\mathrm{w}}(n), \tag{59}$$

with $e_{\mathrm{b}}(n)$ and $e_{\mathrm{w}}(n)$ as in (53)–(54) and $N_1, \ldots, N_K$, $I^{(n)}$, independent, where the N_r are standard normally distributed for $r = 1, \ldots, K$. A comparison on the definition of $\tilde{Q}_n^{\mathrm{b}}$ and $\tilde{Q}_n^{\mathrm{w}}$ with the right hand sides of (53) and (54) and the scaling (52) yields that we have $\mathbb{E}[\tilde{Q}_n^{\mathrm{b}}] = \mathbb{E}[\tilde{Q}_n^{\mathrm{w}}] = 0$ and $\mathrm{Var}(\tilde{Q}_n^{\mathrm{b}}) = \mathrm{Var}(\tilde{Q}_n^{\mathrm{w}}) = 1$ for all $n \geq 2$. Obviously, we also have $\|\tilde{Q}_n^{\mathrm{b}}\|_3, \|\tilde{Q}_n^{\mathrm{w}}\|_3 < \infty$. Hence, ζ_3 distances between X_n, Y_n, $\tilde{Q}_n^{\mathrm{b}}$, $\tilde{Q}_n^{\mathrm{w}}$, and $\mathcal{N}(0,1)$ are finite for all $n \geq 2$. With

$$\begin{aligned}
\tilde{\Delta}(n) &:= \zeta_3^{\vee}((X_n, Y_n), (\mathcal{N}(0,1), \mathcal{N}(0,1))), \\
\tilde{\Delta}_{\mathrm{b}}(n) &:= \zeta_3(X_n, \mathcal{N}(0,1)), \\
\tilde{\Delta}_{\mathrm{w}}(n) &:= \zeta_3(Y_n, \mathcal{N}(0,1))
\end{aligned}$$

we also start with

$$\zeta_3(X_n, \mathcal{N}(0,1)) \leq \zeta_3(X_n, \tilde{Q}_n^{\mathrm{b}}) + \zeta_3(\tilde{Q}_n^{\mathrm{b}}, \mathcal{N}(0,1)).$$

Analogously to the proof of Theorem 6.1 we obtain $\zeta_3(\tilde{Q}_n^{\mathrm{b}}, \mathcal{N}(0,1)) \to 0$ as $n \to \infty$.

The bound for $\zeta_3(X_n, \tilde{Q}_n^{\mathrm{b}})$ is also analogous to the proof of Theorem 6.1, where now is used that ζ_3 is $(3,+)$ ideal instead of $(2,+)$ ideal. This yields

$$\zeta_3(X_n, \tilde{Q}_n^{\mathrm{b}}) \leq \sum_{r=1}^{a+1} \mathbb{E}\left[\left(\frac{\sigma_{\mathrm{b}}(I_r^{(n)})}{\sigma_{\mathrm{b}}(n)}\right)^3 \tilde{\Delta}(I_r^{(n)})\right] + \sum_{r=a+2}^{K} \mathbb{E}\left[\left(\frac{\sigma_{\mathrm{w}}(I_r^{(n)})}{\sigma_{\mathrm{w}}(n)}\right)^3 \tilde{\Delta}(I_r^{(n)})\right].$$

Then we argue as in the previous proof to obtain analogous to (49)

$$\tilde{\Delta}(n) \leq \sum_{r=1}^{a+1} \mathbb{E}\left[\left(\frac{\sigma_{\mathrm{b}}(I_r^{(n)})}{\sigma_{\mathrm{b}}(n)}\right)^3 \tilde{\Delta}(I_r^{(n)})\right] + \sum_{r=a+2}^{K} \mathbb{E}\left[\left(\frac{\sigma_{\mathrm{w}}(I_r^{(n)})}{\sigma_{\mathrm{w}}(n)}\right)^3 \tilde{\Delta}(I_r^{(n)})\right] + o(1).$$

From this estimate we can deduce $\tilde{\Delta}(n) \to 0$ as for $\Delta(n)$ in the proof of Theorem 6.1, where we need to use that from the expansions (51) and Lemma 2.1 we obtain, as $n \to \infty$, that

$$\sum_{r=1}^{a+1} \mathbb{E}\left[\left(\frac{\sigma_{\mathrm{b}}(I_r^{(n)})}{\sigma_{\mathrm{b}}(n)}\right)^3\right] + \sum_{r=a+2}^{K} \mathbb{E}\left[\left(\frac{\sigma_{\mathrm{w}}(I_r^{(n)})}{\sigma_{\mathrm{w}}(n)}\right)^3\right] \to \sum_{r=1}^{K} \mathbb{E}\left[D_r^{3/2}\right] < 1. \tag{60}$$

□

Remarks. (1) Note that the proof of Theorem 6.2 cannot be done in the $\zeta_2^\vee$ metric since the term corresponding to (60) then is

$$\sum_{r=1}^{a+1} \mathbb{E}\left[\left(\frac{\sigma_{\mathrm{b}}(I_r^{(n)})}{\sigma_{\mathrm{b}}(n)}\right)^2\right] + \sum_{r=a+2}^{K} \mathbb{E}\left[\left(\frac{\sigma_{\mathrm{w}}(I_r^{(n)})}{\sigma_{\mathrm{w}}(n)}\right)^2\right] \to \sum_{r=1}^{K} \mathbb{E}\left[D_r\right] = 1,$$

where a limit < 1 is required to obtain $\tilde{\Delta}(n) \to 0$. This is the reason, why we use $\zeta_3^\vee$. It is possible to use $\zeta_s^\vee$ for any $2 < s \leq 3$ leading to the limit $\sum_{r=1}^{K} \mathbb{E}[D_r^s] < 1$.
(2) The case $a - c = (a+b)/2$ differs in the error terms in (50) which then become $\mathrm{O}(\sqrt{n})$. Since the variances in (51) get additional logarithmic factors we still obtain the system (55)–(56) and our proof technique can be applied as well.
(3) The condition $bc > 0$ cannot be dropped. In our approach this would lead to degenerate systems of limit equations that do not identify limit laws. It is known that under $bc = 0$ different asymptotic behavior appears.

6.2 An urn with random replacements

As an example for random entries in the replacement matrix R we consider a simple model with two colors, black and white. In each step when drawing a black ball, a coin is independently tossed to decide whether the black ball is placed back together with another black ball or together with another white ball. The probability for success (a second black ball) is denoted by $0 < \alpha < 1$. Similarly, if a white ball is drawn a coin with probability $0 < \beta < 1$ is tossed to decide whether a second white ball or a black ball is placed back together with the white ball. We denote the replacement matrix by

$$R = \begin{bmatrix} F_\alpha & 1 - F_\alpha \\ 1 - F_\beta & F_\beta \end{bmatrix} \tag{61}$$

where F_α and F_β denote Bernoulli random variables being 1 with probabilities α and β respectively, otherwise 0. This urn model has been introduced in the context of clinical trials and been studied together with generalizations in [28, 29, 27, 26, 19, 3, 4, 12].

The row sums of R in (61) are both deterministically equal to one, hence the urn is balanced. Again, the number of black balls after n draws starting with an initial composition with one black ball is denoted by B_n^{b}, when starting with a white ball by B_n^{w}. According to our approach in section 2 we obtain the recursive equation

$$B_n^{\mathrm{b}} \stackrel{d}{=} B_{I_n}^{\mathrm{b},(1)} + F_\alpha B_{J_n}^{\mathrm{b},(2)} + (1 - F_\alpha)\, B_{J_n}^{\mathrm{w}}, \quad n \geq 1, \tag{62}$$

where $(B_k^{\mathrm{b},(1)})_{0 \leq k < n}$, $(B_k^{\mathrm{b},(2)})_{0 \leq k < n}$, $(B_k^{\mathrm{w}})_{0 \leq k < n}$, F_α and I_n are independent, and $B_k^{\mathrm{b},(1)}$ and $B_k^{\mathrm{b},(2)}$ are distributed as B_k^{b} for $k = 0, \ldots, n-1$, and I_n is uniformly distributed on $\{0, \ldots, n-$

1} while $J_n := n-1-I_n$. (The uniform distribution of I_n follows from the uniform distribution of the number of balls in the $\left[\begin{smallmatrix}1 & 0\\ 0 & 1\end{smallmatrix}\right]$-Pólya urn.) Similarly, we obtain for B_n^{w} that

$$B_n^{\mathrm{w}} \stackrel{d}{=} B_{I_n}^{\mathrm{w},(1)} + F_\beta B_{J_n}^{\mathrm{w},(2)} + (1-F_\beta)\, B_{J_n}^{\mathrm{b}}, \quad n \geq 1, \tag{63}$$

with conditions on independence and identical distributions similar to (62). Together with the initial value $(B_0^{\mathrm{b}}, B_0^{\mathrm{w}}) = (1,0)$ the system of equations (62)–(63) again defines the sequence of pairs of distributions $(\mathcal{L}(B_n^{\mathrm{b}}), \mathcal{L}(B_n^{\mathrm{w}}))_{n\geq 0}$. As a special case of Lemma 2.1 we have

$$(I_n, J_n) \to (U, 1-U) \quad , (n \to \infty), \tag{64}$$

almost surely where U is uniformly distributed on $[0,1]$. Furthermore, we denote for $n \geq 0$

$$\mu_{\mathrm{b}}(n) := \mathbb{E}[B_n^{\mathrm{b}}], \quad \mu_{\mathrm{w}}(n) := \mathbb{E}[B_n^{\mathrm{w}}]. \tag{65}$$

These means have been studied before. We have the following exact formulae:

Lemma 6.3. *For $\mu_{\mathrm{b}}(n)$ and $\mu_{\mathrm{w}}(n)$ as in (65) with $0 < \alpha, \beta < 1$ we have*

$$\mu_{\mathrm{b}}(n) = \frac{1-\beta}{2-\alpha-\beta}\, n + \frac{1-\alpha}{2-\alpha-\beta}\frac{\Gamma(n+\alpha+\beta)}{\Gamma(\alpha+\beta)\Gamma(n+1)} + \frac{1-\beta}{2-\alpha-\beta}, \tag{66}$$

$$\mu_{\mathrm{w}}(n) = \frac{1-\beta}{2-\alpha-\beta}\, n - \frac{1-\beta}{2-\alpha-\beta}\frac{\Gamma(n+\alpha+\beta)}{\Gamma(\alpha+\beta)\Gamma(n+1)} + \frac{1-\beta}{2-\alpha-\beta}. \tag{67}$$

Proof. A proof is based on matrix diagonalization and can elementary be done along the lines of the proof of Lemma 6.7 below. □

As in the example from section 6.1 we have two different types of limit laws, with normal limit for $\alpha + \beta \leq 3/2$ and with non-normal limit for $\alpha+\beta > 3/2$.

The non-normal limit case. We assume that $\lambda := \alpha + \beta - 1 > 1/2$. From Lemma 6.3 we obtain the asymptotic expressions, as $n \to \infty$,

$$\mu_{\mathrm{b}}(n) = c'_{\mathrm{b}} n + d'_{\mathrm{b}} n^\lambda + o(n^\lambda),$$
$$\mu_{\mathrm{w}}(n) = c'_{\mathrm{w}} n + d'_{\mathrm{w}} n^\lambda + o(n^\lambda)$$

with constants

$$c'_{\mathrm{b}} = c'_{\mathrm{w}} = \frac{1-\beta}{1-\lambda}, \quad d'_{\mathrm{b}} = \frac{1-\alpha}{(1-\lambda)\,\Gamma(\lambda+1)}, \quad d'_{\mathrm{w}} = -\frac{1-\beta}{(1-\lambda)\,\Gamma(\lambda+1)}. \tag{68}$$

We use the normalizations $X_0 := Y_0 := 0$ and

$$X_n := \frac{B_n^{\mathrm{b}} - \mu_{\mathrm{b}}(n)}{n^\lambda}, \quad Y_n := \frac{B_n^{\mathrm{w}} - \mu_{\mathrm{w}}(n)}{n^\lambda}, \quad n \geq 1. \tag{69}$$

As in the non-normal case of the example in section 6.1 it is sufficient to use the order of the error term of the mean for the scaling. From (62)–(63) we obtain for $n \geq 1$,

$$X_n \stackrel{d}{=} \left(\frac{I_n}{n}\right)^\lambda X_{I_n}^{(1)} + F_\alpha \left(\frac{J_n}{n}\right)^\lambda X_{J_n}^{(2)} + (1-F_\alpha)\left(\frac{J_n}{n}\right)^\lambda Y_{J_n} + b'_{\mathrm{b}}(n), \tag{70}$$

$$Y_n \stackrel{d}{=} \left(\frac{I_n}{n}\right)^\lambda Y_{I_n}^{(1)} + F_\beta \left(\frac{J_n}{n}\right)^\lambda Y_{J_n}^{(2)} + (1-F_\beta)\left(\frac{J_n}{n}\right)^\lambda X_{J_n} + b'_{\mathrm{w}}(n), \tag{71}$$

with

$$b'_{\mathrm{b}}(n) = d'_{\mathrm{b}}\left(\left(\frac{I_n}{n}\right)^{\lambda} + F_\alpha\left(\frac{J_n}{n}\right)^{\lambda} - 1\right) + d'_{\mathrm{w}}(1-F_\alpha)\left(\frac{J_n}{n}\right)^{\lambda} + o(1),$$

$$b'_{\mathrm{w}}(n) = d'_{\mathrm{w}}\left(\left(\frac{I_n}{n}\right)^{\lambda} + F_\beta\left(\frac{J_n}{n}\right)^{\lambda} - 1\right) + d'_{\mathrm{b}}(1-F_\beta)\left(\frac{J_n}{n}\right)^{\lambda} + o(1),$$

with conditions on independence and identical distributions analogously to (62)–(63). In view of (64) this suggests for limits X and Y of X_n and Y_n that

$$X \stackrel{d}{=} U^{\lambda}X^{(1)} + F_\alpha(1-U)^{\lambda}X^{(2)} + (1-F_\alpha)\,(1-U)^{\lambda}Y^{(1)} + b'_{\mathrm{b}}, \tag{72}$$

$$Y \stackrel{d}{=} U^{\lambda}Y^{(1)} + F_\beta(1-U)^{\lambda}Y^{(2)} + (1-F_\beta)\,(1-U)^{\lambda}X^{(1)} + b'_{\mathrm{w}}, \tag{73}$$

with

$$b'_{\mathrm{b}} = d'_{\mathrm{b}}\left(U^{\lambda} + F_\alpha(1-U)^{\lambda} - 1\right) + d'_{\mathrm{w}}(1-F_\alpha)(1-U)^{\lambda},$$

$$b'_{\mathrm{w}} = d'_{\mathrm{w}}\left(U^{\lambda} + F_\beta(1-U)^{\lambda} - 1\right) + d'_{\mathrm{b}}(1-F_\beta)(1-U)^{\lambda},$$

where $X^{(1)}$, $X^{(2)}$, $Y^{(1)}$, $Y^{(2)}$ and U are independent and $X^{(1)}$, $X^{(2)}$ are distributed as X and $Y^{(1)}$, $Y^{(2)}$ are distributed as Y.

To check that Theorem 5.1 can be applied to the map associated to the system (72)–(73) first note that the form of d'_{b} and d'_{w} in (68) implies $\mathbb{E}[b'_{\mathrm{b}}] = \mathbb{E}[b'_{\mathrm{w}}] = 0$. To check condition (22) note that we have

$$\mathbb{E}\left[U^{2\lambda}\right] + \mathbb{E}\left[F_\alpha(1-U)^{2\lambda}\right] + \mathbb{E}\left[(1-F_\alpha)(1-U)^{2\lambda}\right] = \frac{2}{2\lambda+1} < 1,$$

since $\lambda > 1/2$. Analogously, we have $\mathbb{E}[U^{2\lambda}] + \mathbb{E}[F_\beta(1-U)^{2\lambda}] + \mathbb{E}[(1-F_\beta)(1-U)^{2\lambda}] = 2/(2\lambda+1) < 1$. Together, this verifies condition (22). Hence Theorem 5.1 can be applied and yields a unique fixed-point $(\mathcal{L}(\Lambda'_{\mathrm{b}}), \mathcal{L}(\Lambda'_{\mathrm{w}}))$ in $\mathcal{M}_2^{\mathbb{R}}(0) \times \mathcal{M}_2^{\mathbb{R}}(0)$ to (72)–(73).

Theorem 6.4. *Consider the Pólya urn with random replacement matrix (61) with $\alpha, \beta \in (0,1)$ and $\alpha + \beta > 3/2$ and the normalized numbers X_n and Y_n of black balls as in (69). Furthermore let $(\mathcal{L}(\Lambda'_{\mathrm{b}}), \mathcal{L}(\Lambda'_{\mathrm{w}}))$ denote the in $\mathcal{M}_2^{\mathbb{R}}(0) \times \mathcal{M}_2^{\mathbb{R}}(0)$ unique solution of (72)–(73). Then, as $n \to \infty$,*

$$X_n \xrightarrow{d} \Lambda'_{\mathrm{b}}, \quad Y_n \xrightarrow{d} \Lambda'_{\mathrm{w}}.$$

Proof. Analogously to the proof of Theorem 6.1. □

The normal limit case. Now, we discuss the normal limit case $\lambda := \alpha + \beta - 1 \leq 1/2$. We first assume $\lambda := \alpha + \beta - 1 < 1/2$. The expansions from Lemma 6.3 now imply, as $n \to \infty$

$$\mu_{\mathrm{b}}(n) = c_{\mathrm{b}}n + o(\sqrt{n}), \quad \mu_{\mathrm{w}}(n) = c_{\mathrm{w}}n + o(\sqrt{n}) \tag{74}$$

with c_{b} and c_{w} given in (68). As in the normal limit cases in the examples in section 6.1 we first need asymptotic expressions for the variances. We denote the variances of B_n^{b} and B_n^{w} with $\hat{\sigma}^2_{\mathrm{b}}(n)$ and $\hat{\sigma}^2_{\mathrm{w}}(n)$. These can be obtained from a result of Matthews and Rosenberger [19] for the number of draws of each color as follows:

Lemma 6.5. *We have, as $n \to \infty$,*

$$\hat{\sigma}_{\mathrm{b}}^2(n) = f'_{\mathrm{b}} n + o(n), \quad \hat{\sigma}_{\mathrm{w}}^2(n) = f'_{\mathrm{w}} n + o(n), \tag{75}$$

with

$$f'_{\mathrm{b}} = f'_{\mathrm{w}} = \frac{(1-\alpha)(1-\beta)}{(1-\lambda)^2} \left(\frac{1}{1-2\lambda} - 2\lambda(1+\lambda) \right) > 0.$$

Proof. In [19], for the present urn model, the number N_n of draws within the first n draws in which a black ball is drawn is studied. Starting with one black ball it is established in [19], as $n \to \infty$, that

$$\begin{aligned} \mathbb{E}[N_n] &= \frac{1-\beta}{1-\lambda} n + o(n), \\ \mathrm{Var}(N_n), &= \frac{(1-\alpha)(1-\beta)(3+2\lambda)}{(1-\lambda)^2 (1-2\lambda)} n + o(n). \end{aligned}$$

As each black ball in the urn is either the first ball, or has been added after drawing a black ball and having success in tossing the corresponding coin, or after drawing a white ball and having no success in tossing the coin, we can directly link N_n to B_n^{b}: Denoting the coin flips after drawing black balls by $(F_j^{\mathrm{b}})_{1 \le j \le N_n}$, the coin flips after drawing white balls by $(F_j^{\mathrm{w}})_{1 \le j \le (n-N_n)}$ we have

$$B_n^{\mathrm{b}} = 1 + \sum_{j=1}^{N_n} F_j^{\mathrm{b}} + \sum_{j=1}^{n-N_n} \left(1 - F_j^{\mathrm{w}}\right).$$

Using that all coin flips are independent we obtain from the law of total variance by conditioning on N_n that

$$\begin{aligned} \hat{\sigma}_b^2(n) &= \mathbb{E}\left[\mathrm{Var}\Big(B_n^{\mathrm{b}} \mid N_n\Big)\right] + \mathrm{Var}\Big(\mathbb{E}\left[B_n^{\mathrm{b}} \mid N_n\right]\Big) \\ &= \frac{(1-\alpha)(1-\beta)}{(1-\lambda)^2} \left(\frac{1}{1-2\lambda} - 2\lambda(1+\lambda) \right) n + o(n). \end{aligned}$$

When starting with one white ball, a similar argument gives the corresponding result. ☐

We use the normalizations $X_0 := Y_0 := 0$ and

$$X_n := \frac{B_n^{\mathrm{b}} - \mu_{\mathrm{b}}(n)}{\hat{\sigma}_{\mathrm{b}}(n)}, \quad Y_n := \frac{B_n^{\mathrm{w}} - \mu_{\mathrm{w}}(n)}{\hat{\sigma}_{\mathrm{w}}(n)}, \quad n \ge 1. \tag{76}$$

From the system (62)–(63) we obtain for the scaled quantities X_n, Y_n the system, for $n \ge 1$,

$$\begin{aligned} X_n &\stackrel{d}{=} \frac{\hat{\sigma}_{\mathrm{b}}(I_n)}{\hat{\sigma}_{\mathrm{b}}(n)} X_{I_n}^{(1)} + F_\alpha \frac{\hat{\sigma}_{\mathrm{b}}(J_n)}{\hat{\sigma}_{\mathrm{b}}(n)} X_{J_n}^{(2)} + (1 - F_\alpha) \frac{\hat{\sigma}_{\mathrm{w}}(J_n)}{\hat{\sigma}_{\mathrm{w}}(n)} Y_{J_n} + e'_{\mathrm{b}}(n), \\ Y_n &\stackrel{d}{=} \frac{\hat{\sigma}_{\mathrm{w}}(I_n)}{\hat{\sigma}_{\mathrm{w}}(n)} Y_{I_n}^{(1)} + F_\beta \frac{\hat{\sigma}_{\mathrm{w}}(J_n)}{\hat{\sigma}_{\mathrm{w}}(n)} Y_{J_n}^{(2)} + (1 - F_\beta) \frac{\hat{\sigma}_{\mathrm{b}}(J_n)}{\hat{\sigma}_{\mathrm{b}}(n)} X_{J_n} + e'_{\mathrm{w}}(n), \end{aligned}$$

with conditions on independence and identical distributions analogously to (62)–(63). We have $\|e'_{\mathrm{b}}(n)\|_\infty, \|e'_{\mathrm{w}}(n)\|_\infty \to 0$ since the leading linear terms in the expansions (74) cancel out and the error terms $o(\sqrt{n})$ are asymptotically eliminated by the scaling of order $1/\sqrt{n}$. In view of (64) this suggests for limits X and Y of X_n and Y_n respectively

$$X \stackrel{d}{=} \sqrt{U} X^{(1)} + F_\alpha \sqrt{1-U} X^{(2)} + (1-F_\alpha)\sqrt{1-U} Y^{(1)}, \tag{77}$$

$$Y \stackrel{d}{=} \sqrt{U} Y^{(1)} + F_\beta \sqrt{1-U} Y^{(2)} + (1-F_\beta)\sqrt{1-U} X^{(1)}, \tag{78}$$

where $X^{(1)}$, $X^{(2)}$, $Y^{(1)}$, $Y^{(2)}$ and U are independent and $X^{(1)}$, $X^{(2)}$ are distributed as X and $Y^{(1)}$, $Y^{(2)}$ are distributed as Y. To the map associated to the system (77)–(78) we can apply Theorem 5.2. The conditions (23) and (24) are trivially satisfied. Hence $(\mathcal{N}(0,1), \mathcal{N}(0,1))$ is the unique fixed-point of the associated map in the space $\mathcal{M}_3^{\mathbb{R}}(0,1) \times \mathcal{M}_3^{\mathbb{R}}(0,1)$.

Theorem 6.6. *Consider the Pólya urn with random replacement matrix (61) with $\alpha, \beta \in (0,1)$ and $\alpha + \beta < 3/2$ and the normalized numbers X_n and Y_n of black balls as in (76). Then, as $n \to \infty$,*

$$X_n \xrightarrow{d} \mathcal{N}(0,1), \quad Y_n \xrightarrow{d} \mathcal{N}(0,1). \tag{79}$$

Proof. Analogously to the proof of Theorem 6.2. □

Remark. The case $\alpha + \beta = 3/2$ differs in the error terms in (74) which then become $\mathrm{O}(\sqrt{n})$. Since the variances in (75) get additional logarithmic factors we still obtain the system (77)–(78) and our proof technique does still apply.

6.3 Cyclic urns

We fix an integer $m \geq 2$ and consider an urn with balls of types $1, \ldots, m$. After drawing a ball of type j it is placed back to the urn together with a ball of type $j+1$ if $1 \leq j \leq m-1$ and together with a ball of type 1 if $j = m$. These urn models are called *cyclic urns*. Thus, the replacement matrix of a cyclic urn has the form

$$R = \begin{bmatrix} 0 & 1 & & & 0 \\ & 0 & 1 & & \\ & & 0 & & \\ & & & \ddots & 1 \\ 1 & & & & 0 \end{bmatrix}. \tag{80}$$

We denote by $R_n^{[j]}$ the number of type 1 balls after n draws when initially one ball of type j is contained in the urn. Our recursive approach described above yields the system of recursive distributional equations

$$\begin{aligned} R_n^{[1]} &\stackrel{d}{=} R_{I_n}^{[1]} + R_{J_n}^{[2]}, \qquad (81) \\ R_n^{[2]} &\stackrel{d}{=} R_{I_n}^{[2]} + R_{J_n}^{[3]}, \\ &\vdots \\ R_n^{[m]} &\stackrel{d}{=} R_{I_n}^{[m]} + R_{J_n}^{[1]}, \end{aligned}$$

where, on the right hand sides, I_n and $R_k^{[j]}$ for $j = 1, \ldots, m$, $k = 0, \ldots, n-1$ are independent, I_n uniformly distributed on $\{0, \ldots, n-1\}$ and $J_n = n - 1 - I_n$.

We denote the imaginary unit by i and use the primitive roots of unity

$$\omega := \omega_m := \exp\left(\frac{2\pi \mathrm{i}}{m}\right) =: \lambda + \mathrm{i}\mu \tag{82}$$

with $\lambda, \mu \in \mathbb{R}$. Note that for $2 \leq m \leq 6$ we have $\lambda \leq 1/2$, while for $m \geq 7$ we have $\lambda > 1/2$. Asymptotic expressions for the mean of the $R_n^{[j]}$ can be found (together with further analysis) in [11, 12, 22]. To keep this section self-contained we give an exact formula for later use:

Lemma 6.7. *Let $R_n^{[j]}$ be the number of balls of color 1 after n draws in a cyclic urn with $m \geq 2$ colors, starting with one ball of color j. Then, with $\omega = \omega_m$ as in (82) we have*

$$\mathbb{E}\Big[R_n^{[j]}\Big] = \frac{n+1}{m} + \frac{1}{m} \sum_{k \in \{1,\dots,m\} \setminus \{m/2\}} \frac{\Gamma(n+1+\omega^k)}{\Gamma(n+1)\,\Gamma(\omega^k+1)} \omega^{k(j-1)}. \tag{83}$$

In particular, we have $\mathbb{E}[R_n^{[j]}] = \frac{1}{m} n + \mathrm{O}(1)$ for $m = 2, 3, 4$ and, for $m > 4$, as $n \to \infty$,

$$\mathbb{E}\left[R_n^{[j]}\right] = \frac{1}{m} n + \Re(\kappa_j n^{\mathrm{i}\mu}) n^{\lambda} + o(n^{\lambda}), \qquad \kappa_j := \frac{2\omega^{j-1}}{m\Gamma(\omega+1)}. \tag{84}$$

Proof. Using the system (81), we obtain by conditioning on I_n, for any $1 \leq j \leq m$,

$$\begin{aligned} \mathbb{E}\left[R_n^{[j]}\right] &= \frac{1}{n} \sum_{i=0}^{n-1} \mathbb{E}\left[R_i^{[j]}\right] + \frac{1}{n} \sum_{i=0}^{n-1} \mathbb{E}\left[R_i^{[j+1]}\right] \\ &= \frac{1}{n}\left(\mathbb{E}\left[R_{n-1}^{[j]}\right] + \mathbb{E}\left[R_{n-1}^{[j+1]}\right]\right) + \frac{n-1}{n} \mathbb{E}\left[R_{n-1}^{[j]}\right] \\ &= \mathbb{E}\left[R_{n-1}^{[j]}\right] + \frac{1}{n} \mathbb{E}\left[R_{n-1}^{[j+1]}\right]. \end{aligned}$$

With the column vector $R_n := (R_n^{[1]}, \dots, R_n^{[m]})$, the replacement matrix R in (80) and the identity matrix Id_m this is rewritten as

$$\mathbb{E}\left[R_n\right] = \left(\mathrm{Id}_m + \frac{1}{n} R\right) \mathbb{E}\left[R_{n-1}\right] = \prod_{k=1}^{n} \left(\mathrm{Id}_m + \frac{1}{k} R\right) \mathbb{E}[R_0].$$

The eigenvalues of the replacement matrix are all m-th roots of unity $\omega_\ell := \omega^\ell$, $\ell = 1, \dots, m$, and a possible eigenbasis is $v_\ell := \frac{1}{m}(\omega_\ell^0, \dots, \omega_\ell^{m-1})^t$, $\ell = 1, \dots, m$. Decomposing the mapping induced by R into the projections π_{v_ℓ} onto the respective eigenspaces we obtain

$$\prod_{k=1}^{n} \left(\mathrm{Id}_m + \frac{1}{k} R\right) = \sum_{l=1}^{m} \prod_{k=1}^{n} \left(1 + \frac{1}{k} \omega_l\right) \pi_{v_l} = (n+1)\ \pi_{v_m} + \sum_{l=1}^{m-1} \frac{\Gamma(n+1+\omega_l)}{\Gamma(\omega_l+1)\Gamma(n+1)} \pi_{v_l}.$$

Moreover, $\pi_{v_j}(\mathbb{E}[R_0]) = v_j$ and $v_m = \frac{1}{m}(1, \dots, 1)$, hence the j-th component of the latter display implies (83).

The asymptotic expansion in (84) is now directly read off: Note that the roots of unity come in conjugate pairs $\overline{\omega_\ell} = \omega_{m-\ell}$. If m is even, $\omega_{m/2} = \overline{\omega_{m/2}} = -1$, otherwise only $\omega_m = 1$

is real. Combining pairs of summands for such conjugate pairs and using $\Gamma(\bar{z}) = \overline{\Gamma(z)}$, we obtain the terms

$$\frac{\Gamma(n+1+\omega_\ell)\,\omega_\ell^{j-1}}{\Gamma(n+1)\,\Gamma(\omega_\ell+1)} + \frac{\Gamma(n+1+\bar{\omega_\ell})\,\bar{\omega_\ell}^{j-1}}{\Gamma(n+1)\,\Gamma(\bar{\omega_\ell}+1)} = 2\,\Re\!\left(\frac{\omega_\ell^{j-1}\,\Gamma(n+1+\omega_\ell)}{\Gamma(\omega_\ell+1)\,\Gamma(n+1)}\right).$$

By Stirling approximation the asymptotic growth order of the latter term is $\Re(n^{\omega_\ell})$, hence the dominant asymptotic term is for the conjugate pair with largest real part, ω and ω_{m-1}. This implies (84) for $m > 4$. For $m = 3, 4$ the periodic term is $o(1)$ respectively $\mathrm{O}(1)$, for $m = 2$ there is no periodic fluctuation. □

We do not discuss limit laws for the cases $2 \le m \le 6$ in detail. They lead to asymptotic normality as has been shown with different proofs in Janson [11] and Janson [12, Example 7.9]. These cases can be covered by our approach similarly to the normal cases in sections 6.1 and 6.2. For $2 \le m \le 6$, the system of limit equations is

$$\begin{aligned}
X^{[1]} &\stackrel{d}{=} \sqrt{U}X^{[1]} + \sqrt{1-U}X^{[2]},\\
X^{[2]} &\stackrel{d}{=} \sqrt{U}X^{[2]} + \sqrt{1-U}X^{[3]},\\
&\vdots\\
X^{[m]} &\stackrel{d}{=} \sqrt{U}X^{[m]} + \sqrt{1-U}X^{[1]},
\end{aligned}$$

and Theorem 5.2 applies.

We now assume $m \ge 7$. In particular, we have the asymptotic expansion (84) of the mean of the $R_n^{[j]}$ with $\lambda > 1/2$. We define the normalizations

$$X_n^{[j]} := \frac{R_n^{[j]} - \frac{1}{m}n}{n^\lambda}. \tag{85}$$

Hence, we obtain for the $X_n^{[j]}$ the system

$$\begin{aligned}
X_n^{[1]} &\stackrel{d}{=} \left(\frac{I_n}{n}\right)^\lambda X_{I_n}^{[1]} + \left(\frac{J_n}{n}\right)^\lambda X_{J_n}^{[2]} - \frac{1}{mn^\lambda},\\
X_n^{[2]} &\stackrel{d}{=} \left(\frac{I_n}{n}\right)^\lambda X_{I_n}^{[2]} + \left(\frac{J_n}{n}\right)^\lambda X_{J_n}^{[3]} - \frac{1}{mn^\lambda},\\
&\vdots\\
X_n^{[m]} &\stackrel{d}{=} \left(\frac{I_n}{n}\right)^\lambda X_{I_n}^{[m]} + \left(\frac{J_n}{n}\right)^\lambda X_{J_n}^{[1]} - \frac{1}{mn^\lambda},
\end{aligned}$$

where, on the right hand sides, I_n and $X_k^{[j]}$ for $j = 1, \dots, m$, $k = 0, \dots, n-1$ are independent. To describe the asymptotic periodic behavior of the distributions of the $X_n^{[j]}$, we use the following related system of limit equations:

$$\begin{aligned}
X^{[1]} &\stackrel{d}{=} U^\omega X^{[1]} + (1-U)^\omega X^{[2]},\\
X^{[2]} &\stackrel{d}{=} U^\omega X^{[2]} + (1-U)^\omega X^{[3]},\\
&\vdots\\
X^{[m]} &\stackrel{d}{=} U^\omega X^{[m]} + (1-U)^\omega X^{[1]}.
\end{aligned}$$

Since ω is complex this now has to be considered as a system to solve for distributions $\mathcal{L}(X^{[1]}), \ldots, \mathcal{L}(X^{[m]})$ on the complex plane $\mathbb{C}$. The corresponding map $\bar{T}$ is a special case of T' in (21):

$$\begin{aligned}\bar{T} : \mathcal{M}^{\mathbb{C},\times m} &\to \mathcal{M}^{\mathbb{C},\times m}\\ (\mu_1, \ldots, \mu_m) &\mapsto (\bar{T}_1(\mu_1, \ldots, \mu_m), \ldots, \bar{T}_m(\mu_1, \ldots, \mu_m))\\ \bar{T}_j(\mu_1, \ldots, \mu_m) &:= \mathcal{L}\left(U^\omega V^{[j]} + (1-U)^\omega V^{[j+1]}\right) \qquad (86)\end{aligned}$$

for $j = 1, \ldots, m$ with $U, V^{[1]}, \ldots, V^{[m+1]}$ independent, U uniformly distributed on $[0, 1]$ and $\mathcal{L}(V^{[j]}) = \mu_j$ for $j = 1, \ldots, m$ and $\mathcal{L}(V^{[m+1]}) = \mu_1$.

Lemma 6.8. *Let $m \geq 7$. The restriction of $\bar{T}$ to $\mathcal{M}_2^{\mathbb{C}}(\kappa_1) \times \cdots \times \mathcal{M}_2^{\mathbb{C}}(\kappa_m)$ has a unique fixed-point.*

Proof. We verify the conditions of Theorem 5.3: First note that condition (25) for our $\bar{T}$ in (86) is

$$\mathbb{E}\left[U^\omega\right]\kappa_j + \mathbb{E}\left[(1-U)^\omega\right]\kappa_{j+1} = \kappa_j, \quad j = 1, \ldots, m, \qquad (87)$$

with $\kappa_{m+1} := \kappa_1$. Since $\mathbb{E}[U^\omega] = \mathbb{E}[(1-U)^\omega] = (1+\omega)^{-1}$ and $\kappa_{j+1} = \omega\kappa_j$ we find that (87) is satisfied. Condition (26) for our $\bar{T}$ is

$$\mathbb{E}\left[\left|U^{2\omega}\right|\right] + \mathbb{E}\left[\left|(1-U)^{2\omega}\right|\right] < 1.$$

Since $m \geq 7$, we have $\lambda > 1/2$, thus $\mathbb{E}[|U^{2\omega}|] + \mathbb{E}[|(1-U)^{2\omega}|] = 2/(1+2\lambda) < 1$. Hence Theorem 5.3 applies and implies the assertion. $\square$

The fixed-point in Lemma 6.8 has a particularly simple structure as follows:

Lemma 6.9. *Let $m \geq 7$ and $(\mathcal{L}(\Lambda^{[1]}), \ldots, \mathcal{L}(\Lambda^{[m]}))$ be the unique fixed-point in Lemma 6.8. Furthermore let $\mathcal{L}(\Lambda)$ be the (unique) fixed-point of*

$$X \stackrel{d}{=} U^\omega X + \omega(1-U)^\omega X' \quad \text{in} \quad \mathcal{M}_2^{\mathbb{C}}\left(\frac{2}{m\Gamma(\omega+1)}\right), \qquad (88)$$

where X, X' and U are independent, U is uniformly distributed on $[0, 1]$ and X and X' have identical distributions. Then we have

$$\Lambda^{[j]} \stackrel{d}{=} \omega^{j-1}\Lambda, \quad j = 1, \ldots, m.$$

Proof. We abbreviate $\gamma := 2/(m\Gamma(\omega+1))$. For X, X' and U independent, U uniformly on $[0, 1]$ distributed, X and X' identically distributed with $\mathbb{E}X = \gamma$ we have

$$\mathbb{E}\left[U^\omega X + \omega(1-U)^\omega X'\right] = \frac{1}{1+\omega}(\gamma + \omega\gamma) = \gamma,$$

hence the map of probability measures on $\mathbb{C}$ associated to (88) maps $\mathcal{M}_2^{\mathbb{C}}(\gamma)$ into itself. The argument of the proof of Theorem 5.3 implies that this map is a contraction on $(\mathcal{M}_2^{\mathbb{C}}(\gamma), \ell_2)$. Hence it has a unique fixed point $\mathcal{L}(\Lambda)$. We have

$$(\mathcal{L}(\Lambda), \mathcal{L}(\omega\Lambda), \ldots, \mathcal{L}(\omega^{m-1}\Lambda)) \in \mathcal{M}_2^{\mathbb{C}}(\kappa_1) \times \cdots \times \mathcal{M}_2^{\mathbb{C}}(\kappa_m)$$

and, by plugging into (86), we find that this vector is a fixed-point of $\bar{T}$. Since, by Lemma 6.8, there is only one fixed-point of $\bar{T}$ in $\mathcal{M}_2^{\mathbb{C}}(\kappa_1) \times \cdots \times \mathcal{M}_2^{\mathbb{C}}(\kappa_m)$ the assertion follows. $\square$

The asymptotic periodic behavior in the following theorem has already been shown almost surely by martingale methods in [22, Section 4.2], see also [12, Theorem 3.24]. Our contraction approach adds the characterization of $\mathcal{L}(\Lambda)$ as the fixed-point in (88). The proof is based on the complex version of the ℓ_2 metric and resembles ideas from Fill and Kapur [9], see also [13, Theorem 5.3].

Theorem 6.10. *Let $m \geq 7$ and $X_n^{[j]}$ as in (85) and $\mathcal{L}(\Lambda)$ the unique fixed-point in Lemma 6.9. Then, for all $j = 1, \dots, m$, we have*

$$\ell_2 \left(X_n^{[j]}, \Re \left(e^{\mathrm{i}\left(\mu \ln(n) + 2\pi \frac{j-1}{m}\right)} \Lambda \right) \right) \to 0 \quad (n \to \infty). \tag{89}$$

Proof. Let $\Lambda^{[1]}, \dots, \Lambda^{[m]}$ be independent random variables such that $(\mathcal{L}(\Lambda^{[1]}), \dots, \mathcal{L}(\Lambda^{[m]}))$ is the unique fixed-point as in Lemma 6.8. Set $\Lambda^{[m+1]} := \Lambda^{[1]}$. Note that for the random variable within the real part in (89) with Lemma 6.9 we have

$$e^{\mathrm{i}\left(\mu \ln(n) + 2\pi \frac{j-1}{m}\right)} \Lambda = n^{\mathrm{i}\mu} \omega^{j-1} \Lambda \stackrel{d}{=} n^{\mathrm{i}\mu} \Lambda^{[j]}. \tag{90}$$

The fixed-point property of the $\Lambda^{[j]}$ implies

$$\Re \left(n^{\mathrm{i}\mu} \Lambda^{[j]} \right) \stackrel{d}{=} \Re \left(n^{\mathrm{i}\mu} U^{\omega} \Lambda^{[j]} \right) + \Re \left(n^{\mathrm{i}\mu} (1-U)^{\omega} \Lambda^{[j+1]} \right) \tag{91}$$

for all $j = 1, \dots, m$ and $n \geq 0$. We denote

$$\Delta_j(n) := \ell_2 \left(X_n^{[j]}, \Re \left(n^{\mathrm{i}\mu} \Lambda^{[j]} \right) \right) \tag{92}$$

and set $\Delta_{m+1}(n) := \Delta_1(n)$. Now, we assume that the $X_n^{[j]}$, $\Lambda^{[j]}$, $n \geq 1$, $1 \leq j \leq m$, I_n, U appearing in (85) and (86) are defined on one probability space such that $(X_n^{[j]}, \Re(n^{\mathrm{i}\mu} \Lambda^{[j]}))$ are optimal ℓ_2-couplings for all $n \geq 0$ and all $1 \leq j \leq m$ and such that $I_n = \lfloor nU \rfloor$. Then we have

$$\begin{aligned}
&\Delta_j(n) \\
&= \ell_2 \left(\left(\frac{I_n}{n} \right)^{\lambda} X_{I_n}^{[j]} + \left(\frac{J_n}{n} \right)^{\lambda} X_{J_n}^{[j+1]} - \frac{1}{mn^{\lambda}}, \Re \left(n^{\mathrm{i}\mu} U^{\omega} \Lambda^{[j]} \right) + \Re \left(n^{\mathrm{i}\mu} (1-U)^{\omega} \Lambda^{[j+1]} \right) \right) \\
&\leq \left\| \left\{ \left(\frac{I_n}{n} \right)^{\lambda} X_{I_n}^{[j]} - \Re \left(\frac{I_n^{\omega}}{n^{\lambda}} \Lambda^{[j]} \right) \right\} + \left\{ \left(\frac{J_n}{n} \right)^{\lambda} X_{J_n}^{[j+1]} - \Re \left(\frac{J_n^{\omega}}{n^{\lambda}} \Lambda^{[j+1]} \right) \right\} \right\|_2 \\
&\quad + \left\| \Re \left(\frac{I_n^{\omega}}{n^{\lambda}} \Lambda^{[j]} \right) - \Re \left(n^{\mathrm{i}\mu} U^{\omega} \Lambda^{[j]} \right) \right\|_2 + \left\| \Re \left(\frac{J_n^{\omega}}{n^{\lambda}} \Lambda^{[j+1]} \right) - \Re \left(n^{\mathrm{i}\mu} U^{\omega} \Lambda^{[j+1]} \right) \right\|_2 + \frac{1}{mn^{\lambda}} \\
&=: S_1 + S_2 + S_3 + \frac{1}{mn^{\lambda}}.
\end{aligned} \tag{93}$$

First note that the summands S_2 and S_3 tend to zero: We have $(I_n/n)^{\omega} \to U^{\omega}$ almost surely by $I_n = \lfloor nU \rfloor$. Since $\Lambda^{[j]}$ and $\Lambda^{[j+1]}$ have finite second moments we can apply dominated convergence to obtain $S_2, S_3 \to 0$ as $n \to \infty$.

For the estimate of the first summand S_1 we abbreviate

$$W_n^{[j]} := \left(\frac{I_n}{n} \right)^{\lambda} X_{I_n}^{[j]} - \Re \left(\frac{I_n^{\omega}}{n^{\lambda}} \Lambda^{[j]} \right), \qquad W_n^{[j+1]} := \left(\frac{J_n}{n} \right)^{\lambda} X_{J_n}^{[j+1]} - \Re \left(\frac{J_n^{\omega}}{n^{\lambda}} \Lambda^{[j+1]} \right).$$

Then we have

$$S_1^2 = \mathbb{E}\left[(W_n^{[j]})^2\right] + \mathbb{E}\left[(W_n^{[j+1]})^2\right] + 2\mathbb{E}\left[W_n^{[j]}W_n^{[j+1]}\right] \tag{94}$$

Conditioning on I_n and using that $(X_k^{[j]}, \Re(k^{\mathrm{i}\mu}\Lambda^{[j]}))$ are optimal ℓ_2-couplings we obtain

$$\begin{aligned}\mathbb{E}\left[(W_n^{[j]})^2\right] &= \sum_{k=0}^{n-1} \frac{1}{n}\mathbb{E}\left[\left\{\left(\frac{k}{n}\right)^{\lambda} X_k^{[j]} - \Re\left(\frac{k^{\lambda}k^{\mathrm{i}\mu}}{n^{\lambda}}\Lambda^{[j]}\right)\right\}^2\right] \\ &= \sum_{k=0}^{n-1} \frac{1}{n}\left(\frac{k}{n}\right)^{2\lambda}\mathbb{E}\left[\left\{X_k^{[j]} - \Re\left(k^{\mathrm{i}\mu}\Lambda^{[j]}\right)\right\}^2\right] \\ &= \sum_{k=0}^{n-1} \frac{1}{n}\left(\frac{k}{n}\right)^{2\lambda}\Delta_j^2(k) \\ &= \mathbb{E}\left[\left(\frac{I_n}{n}\right)^{2\lambda}\Delta_j^2(I_n)\right].\end{aligned}$$

Analogously, we have

$$\mathbb{E}[(W_n^{[j+1]})^2] = \mathbb{E}\left[\left(\frac{J_n}{n}\right)^{2\lambda}\Delta_{j+1}^2(J_n)\right].$$

To bound the mixed term in (94) note that by the expansion (84) and the normalization (85) we have $\mathbb{E}[X_n^{[j]}] = \Re(\kappa_j n^{\mathrm{i}\mu}) + r_j(n)$ with $r_j(n) \to 0$ as $n \to \infty$ for all $j = 1, \dots, m$. In particular, we have $\|r_j\|_\infty < \infty$. Together with $\mathbb{E}[\Lambda^{[j]}] = \kappa_j$ this implies $\mathbb{E}[W_n^{[j]}] = \mathbb{E}[(I_n/n)^{\lambda} r_j(I_n)]$ and

$$\mathbb{E}\left[W_n^{[j]}W_n^{[j+1]}\right] = \mathbb{E}\left[\left(\frac{I_n}{n}\frac{J_n}{n}\right)^{\lambda} r_j(I_n) r_{j+1}(J_n)\right]. \tag{95}$$

To show that the latter term tends to zero let $\varepsilon > 0$. Then there exists $k_0 \in \mathbb{N}$ such that $r_j(k) < \varepsilon$, $r_{j+1}(k) < \varepsilon$ for all $k \geq k_0$. For all $n > 2k_0$ we obtain, by considering the event $\{k_0 \leq I_n \leq n-1-k_0\}$ and its complement,

$$\mathbb{E}\left[W_n^{[j]}W_n^{[j+1]}\right] \leq \frac{2k_0}{n}\|r_j\|_\infty \|r_{j+1}\|_\infty + \varepsilon^2.$$

Hence, we obtain that the mixed term (95) tends to zero.

Altogether, we obtain from (93) as $n \to \infty$ that

$$\begin{aligned}\Delta_j(n) &\leq \left\{\mathbb{E}\left[\left(\frac{I_n}{n}\right)^{2\lambda}\Delta_j^2(I_n)\right] + \mathbb{E}\left[\left(\frac{J_n}{n}\right)^{2\lambda}\Delta_{j+1}^2(J_n)\right] + o(1)\right\}^{1/2} + o(1) \\ &\leq \left\{2\mathbb{E}\left[\left(\frac{I_n}{n}\right)^{2\lambda}\Delta^2(I_n)\right] + o(1)\right\}^{1/2} + o(1),\end{aligned}$$

for all $j = 1, \dots, m$, where

$$\Delta(n) := \max_{1 \le j \le m} \Delta_j(n).$$

Hence, we have

$$\Delta(n) \le \left\{ 2\mathbb{E}\left[\left(\frac{I_n}{n} \right)^{2\lambda} \Delta^2(I_n) \right] + o(1) \right\}^{1/2} + o(1). \tag{96}$$

Now, we obtain $\Delta(n) \to 0$ as in the proof of Theorem 6.1: First from (96) we obtain with $I_n/n \to U$ almost surely that

$$\begin{aligned} \Delta(n) &\le \left\{ 2\mathbb{E}\left[\left(\frac{I_n}{n} \right)^{2\lambda} \right] \max_{0 \le k \le n-1} \Delta^2(k) + o(1) \right\}^{1/2} + o(1) \\ &\le \left\{ \left(\frac{2}{1+2\lambda} + o(1) \right) \max_{0 \le k \le n-1} \Delta^2(k) + o(1) \right\}^{1/2} + o(1). \end{aligned}$$

Since $\lambda > 1/2$ this implies that the sequence $(\Delta(n))_{n \ge 0}$ is bounded. We denote $\eta := \sup_{n \ge 0} \Delta(n)$ and $\xi := \limsup_{n \to \infty} \Delta(n)$. For any $\varepsilon > 0$ there exists an $n_0 \ge 0$ such that $\Delta(n) \le \xi + \varepsilon$ for all $n \ge n_0$. Hence, from (96) we obtain

$$\Delta(n) \le \left\{ 2\mathbb{E}\left[\mathbf{1}_{\{I_n < n_0\}} \left(\frac{I_n}{n} \right)^{2\lambda} \right] \eta^2 + 2\mathbb{E}\left[\mathbf{1}_{\{I_n \ge n_0\}} \left(\frac{I_n}{n} \right)^{2\lambda} \right] (\xi + \varepsilon)^2 + o(1) \right\}^{1/2} + o(1).$$

With $n \to \infty$ this implies

$$\xi \le \sqrt{\frac{2}{1+2\lambda}} (\xi + \varepsilon).$$

Since $\sqrt{2/(1+2\lambda)} < 1$ and $\varepsilon > 0$ is arbitrary this implies $\xi = 0$. □

References

[1] Athreya, K.B. and Karlin, S. (1968) Embedding of urn schemes into continuous time Markov branching processes and related limit theorems. *Ann. Math. Statist.* **39**, 1801–1817.

[2] Bagchi, A. and Pal, A.K. (1985) Asymptotic Normality in the Generalized PolyaEggenberger Urn Model, with an Application to Computer Data Structures. *SIAM J. Algebraic Discrete Methods* **6**, 394–405.

[3] Bai, Z.D. and Hu, F. (1999). Asymptotic theorems for urn models with nonhomogeneous generating matrices. *Stochastic Process. Appl.* **80**, 87–101.

[4] Bai, Z.D., Hu, F. and Zhang, L.-X. (2002). Gaussian approximation theorems for urn models and their applications. *Ann. Appl. Probab.* **12**, 1149–1173.

[5] Bickel, P. J. and Freedman, P. A. (1981) Some asymptotic theory for the bootstrap. *Ann. Statist.* **9**, 1196–1217.

[6] Chauvin, B., Pouyanne, N. and Sahnoun, R. (2011) Limit distributions for large Pólya urns. *Ann. Appl. Probab.* **21**, 1–32.

[7] Chauvin, B., Liu, Q. and Pouyanne, N. (2012) Limit distributions for multitype branching processes of m-ary search trees. *Ann. IHP*, to appear.

[8] Drmota, M., Janson, S. and Neininger, R. (2008) A functional limit theorem for the profile of search trees. *Ann. Appl. Probab.* **18**, 288–333.

[9] Fill, J.A. and Kapur, N. (2004) The Space Requirement of m-ary Search Trees: Distributional Asymptotics for $m \geq 27$. Invited paper, Proceedings of the 7th Iranian Statistical Conference, 2004. Available via `http://www.ams.jhu.edu/~fill/papers/periodic.pdf`

[10] Flajolet, P., Gabarró, J. and Pekari, H. (2005) Analytic urns. *Ann. Probab.* **33**, 1200–1233.

[11] Janson, S. (1983) Limit theorems for certain branching random walks on compact groups and homogeneous spaces. *Ann. Probab.* **11**, 909–930.

[12] Janson, S. (2004) Functional limit theorem for multitype branching processes and generalized Pólya urns. *Stochastic Process. Appl.* **110**, 177–245.

[13] Janson, S. and Neininger, R. (2008) The size of random fragmentation trees. *Probab. Theory Related Fields* **142**, 399–442.

[14] Johnson, N.L. and Kotz, S. (1977) Urn models and their application. An approach to modern discrete probability theory. Wiley Series in Probability and Mathematical Statistics. *John Wiley & Sons, New York-London-Sydney.*

[15] Kaijser, S. and Janson, S. (2012) Higher moments of Banach space valued random variables. Preprint available via `http://arxiv.org/abs/1208.4272`

[16] Kotz, S., Mahmoud, H.M. and Robert, P. (2000) On generalized Pólya urn models. *Statist. Probab. Lett.* **49**, 163–173.

[17] Leckey, K., Neininger, R. and Szpankowski, W. (2013) Towards More Realistic Probabilistic Models for Data Structures: The External Path Length in Tries under the Markov Model. *Proceedings of ACM-SIAM Symposium on Discrete Algorithms (SODA)*, 877–886.

[18] Mahmoud, H.M. (2009) Pólya urn models. Texts in Statistical Science Series. *CRC Press, Boca Raton, FL.*

[19] Matthews, P.C. and Rosenberger, W.F. (1997) Variance in randomized play-the-winner clinical trials. *Statistics & Probability Letters* **35**, 233–240.

[20] Neininger, R. and Rüschendorf, L. (2004) A general limit theorem for recursive algorithms and combinatorial structures. *Ann. Appl. Probab.* **14**, 378–418.

[21] Neininger, R. and Sulzbach, H. (2012) On a functional contraction method. Preprint available via `http://arxiv.org/abs/1202.1370`

[22] Pouyanne, N. (2005) Classification of large Pólya-Eggenberger urns with regard to their asymptotics. *2005 International Conference on Analysis of Algorithms*, 275–285 (electronic), Discrete Math. Theor. Comput. Sci. Proc., AD, *Assoc. Discrete Math. Theor. Comput. Sci., Nancy.*

[23] Pouyanne, N. (2008) An algebraic approach to Pólya processes. *Ann. Inst. Henri PoincarṔrobab. Stat.* **44**, 293–323.

[24] Rachev, S.T. and Rüschendorf, L. (1995) Probability metrics and recursive algorithms. *Adv. in Appl. Probab.* **27**, 770–799.

[25] Rösler, U. (1991) A limit theorem for "Quicksort". *RAIRO Inform. Théor. Appl.* **25**, 85–100.

[26] Smythe, R.T. (1996). Central limit theorems for urn models. *Stochastic Process. Appl.* **65**, 115–137.

[27] Smythe, R.T. and Rosenberger, W.F. (1995). Play-the-winner designs, generalized Pṕlya urns, and Markov branching processes. In: Flournoy, N., Rosenberger, W.F. (Eds.), Adaptive Designs, IMS Lecture Notes Monograph Ser., Vol. 25. Inst. Math. Statist., Hayward, CA, pp. 1322.

[28] Wei, L.J. and Durham, S. (1978). The randomized play-the-winner rule in medical trials. *J. Amer. Statist. Assoc.* **73**, 840–843.

[29] Wei, L.J., Smythe, R.T., Lin, D.Y. and Park, T.S. (1990). Statistical inference with data-dependent treatment allocation rules. *J. Amer. Statist. Assoc.* **85**, 156–162.

[30] Zolotarev, V. M. (1976) Approximation of the distributions of sums of independent random variables with values in infinite-dimensional spaces. (Russian.) *Teor. Veroyatnost. i Primenen.* **21**, 741–758. Erratum *ibid* **22** (1977), 901. English transl. *Theory Probab. Appl.* **21**, 721–737; *ibid.* **22**, 881.

[31] Zolotarev, V. M. (1977) Ideal metrics in the problem of approximating the distributions of sums of independent random variables. (Russian.) *Teor. Veroyatnost. i Primenen.* **22**, 449–465. English transl. *Theory Probab. Appl.* **22**, 433–449.